



[Go to Product page](#)

Datasheet for ABIN657401
anti-CCDC117 antibody (C-Term)

2 Images

Overview

Quantity:	400 µL
Target:	CCDC117
Binding Specificity:	AA 143-171, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This CCDC117 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 143-171 amino acids from the C-terminal region of human CCDC117.
Clone:	RB33331
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CCDC117
Alternative Name:	CCDC117 (CCDC117 Products)
Background:	The specific function of the protein remains unknown.
Molecular Weight:	30541

Target Details

Gene ID: 150275

NCBI Accession: [NP_775781](#)

UniProt: [Q8IWD4](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: CCD117 Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C.


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. CCD Antibody (C-term) (ABIN657401 and ABIN2846441) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CCD Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Anti-CCD Antibody (C-term) at 1:1000 dilution + COL whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.