

Datasheet for ABIN6579823
anti-CD83 antibody (APC)[Go to Product page](#)

1 Image

Overview

Quantity:	100 tests
Target:	CD83
Reactivity:	Human, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD83 antibody is conjugated to APC
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Human CD83-transfected Cos cells
Clone:	HB15e
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody HB15e recognizes an extracellular epitope of CD83, a 40-45 kDa type I glycoprotein expressed on mature dendritic cells.
Cross-Reactivity (Details):	Human, Non-Human Primates
Purification:	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Target Details

Target:	CD83
---------	------

Target Details


Alternative Name:	CD83 (CD83 Products)
Background:	CD83 Molecule,CD83 is a 40-45 kDa heavily glycosylated type I cell surface glycoprotein of immunoglobulin family. It is expressed on the surface of mature dendritic cells, Langerhans cells in the skin, and interdigitating reticulum cells in the lymphoid tissues. Low expression of CD83 has been reported in activated T and B cells. Cytoplasmic expression of CD83 can be detected also in monocytes and macrophages. CD83 is involved in modulation of antigen presentation. Soluble CD83 has immunoregulatory functions, it is able to down-regulate dendritic cell maturation and stimulation of T cells. In the developing immune system, release of soluble CD83 from dendritic cells upon stimulation by gram-positive or gram-negative bacteria has anti-allergic effect. Herpes simplex virus, on the other hand, causes CD83 degradation in mature dendritic cells.,BL11, HB15
Gene ID:	9308
UniProt:	Q01151

Application Details

Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µL reagent / 100 µL of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Comment:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Restrictions:	For Research Use only

Handling

Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.



Flow Cytometry

Image 1. Flow cytometry analysis (surface staining) of CD83 in IFN alpha-activated human peripheral blood cells with anti-CD83 (HB15e) APC.