

Datasheet for ABIN6579828

anti-CD68 antibody (APC)

2 Images



Go to Product page

Overview


| Quantity: | 100 tests |
|--------------|---|
| Target: | CD68 |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This CD68 antibody is conjugated to APC |
| Application: | Flow Cytometry (FACS) |


Product Details

| Purpose: | Anti-Hu CD68 APC |
|-----------------------------|---|
| Immunogen: | Lysosomal contents of lung macrophages |
| Clone: | Y1-82A |
| Isotype: | lgG2b |
| Specificity: | The mouse monoclonal antibody Y1/82A recognizes CD68 (LAMP4), a 110 kDa glycoprotein expressed mainly in cytoplasmic granules of monocytes/macrophages, granulocytes, and dendritic cells. |
| Cross-Reactivity (Details): | Human |
| Purification: | Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |

Target Details

| Target: | CD68 |
|---------------------|--|
| Alternative Name: | CD68 (CD68 Products) |
| Background: | CD68 Molecule,CD68 (also known as LAMP4 or SCARD1) is a 110 kDa type I transmembrane glycoprotein of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family and the scavenger receptor family. Although CD68 primarily localizes to lysosomes and endosomes, its fraction circulates to the cell surface. By the heavily glycosylated extracellular domain CD68 binds to tissue- and organ-specific lectins or selectins. It is expressed mainly in cytoplasmic granules of monocytes/macrophages, granulocytes, and dendritic cells, but also e.g. in a proportion of epithelial tumours (diagnosis of poorly differentiated neoplasms).,GP110, LAMP4, SCARD1 |
| Gene ID: | 968 |
| UniProt: | P34810 |
| Application Details | |
| Application Notes: | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ L reagent / 100 μ L of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient fo 100 tests. Extracellular and intracellular staining. |
| Restrictions: | For Research Use only |
| Handling | |
| Buffer: | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze. |





Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD68 (Y1/82A) APC antibody (10 μ L reagent / 100 μ L of peripheral whole blood).

Flow Cytometry

Image 2. Separation of human monocytes (red-filled) from human CD68 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD68 (Y1/82A) APC antibody (10 µL reagent / 100 µL of peripheral whole blood).