

Datasheet for ABIN6654166  
**anti-IGFBP2 antibody (AA 35-304)**

5 Images



[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	IGFBP2
Binding Specificity:	AA 35-304
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGFBP2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	Rat IGFBP2 recombinant protein (amino acids E35-Q304) was used as the immunogen for the IGFBP2 antibody.
Isotype:	IgG
Purification:	Antigen affinity purified

## Target Details

Target:	IGFBP2
Alternative Name:	IGFBP2 ( <a href="#">IGFBP2 Products</a> )
Background:	The superfamily of insulin-like growth factor (IGF) binding proteins include the six high-affinity IGF binding proteins (IGFBP) and at least four additional low-affinity binding proteins referred to as IGFBP related proteins (IGFBP-rP). All IGFBP superfamily members are cysteine-rich proteins

## Target Details

---

with conserved cysteine residues, which are clustered in the amino- and carboxy-terminal thirds of the molecule. IGFBPs modulate the biological activities of IGF proteins. Some IGFBPs may also have intrinsic bioactivity that is independent of their ability to bind IGF proteins. Post-translational modifications of IGFBPs, including glycosylation, phosphorylation and proteolysis, have been shown to modify the affinities of the binding proteins to IGF. Human IGFBP-2 cDNA encodes a 328 amino acid (aa) residue precursor protein with a putative 39 aa residue signal peptide that is processed to generate the 289 aa residue mature protein. IGFBP-2 contains an integrin receptor recognition sequence (RGD sequence) but lacks potential N-linked glycosylation sites. During development, IGFBP-2 is expressed in a number of tissues. The highest expression level is found in the central nervous system. In adults, high expression levels are also detected in the central nervous system and in a number of reproductive tissues. IGFBP-2 binds preferentially to IGF II, exhibiting a 2-10 fold higher affinity for IGF II than for IGF I.

---

UniProt: [P12843](#)

---

Pathways: [Myometrial Relaxation and Contraction](#), [Growth Factor Binding](#), [Activated T Cell Proliferation](#)

## Application Details

---

Application Notes: Optimal dilution of the IGFBP2 antibody should be determined by the researcher.\. Western blot: 0.5-1 µg/mL, Immunohistochemistry (FFPE): 1-2 µg/mL, Direct ELISA: 0.1-0.5 µg/mL (recombinant rat protein)

---

Restrictions: For Research Use only

## Handling

---


Buffer: 0.5 mg/mL if reconstituted with 0.2 mL sterile DI water

---

Storage: -20 °C


---

Storage Comment: After reconstitution, the IGFBP2 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.




#### Immunohistochemistry

**Image 1.** IHC testing of FFPE mouse kidney tissue with IGFBP2 antibody at 2ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



#### Immunohistochemistry

**Image 2.** IHC testing of FFPE mouse brain tissue with IGFBP2 antibody at 2ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



#### Immunohistochemistry

**Image 3.** IHC testing of FFPE rat brain tissue with IGFBP2 antibody at 2ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN6654166.