

Datasheet for ABIN6654203  
**anti-ALK antibody (AA 200-335)**



[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	ALK
Binding Specificity:	AA 200-335
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This ALK antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	A portion of amino acids 200-335 from the human protein was used as the immunogen for this ALK antibody.
Clone:	ALK-1504
Isotype:	IgG2b kappa
Purification:	Purified
Purity:	Protein G affinity

## Target Details

Target:	ALK
Alternative Name:	ALK / Anaplastic Lymphoma Kinase ( <a href="#">ALK Products</a> )

## Target Details

---

**Background:** The wild-type anaplastic lymphoma kinase (ALK) protein is a 200 kDa transmembrane receptor tyrosine kinase. Its expression is restricted to a few scattered cells in the nervous system (some glial cells and neurons, and a few endothelial cells and pericytes. The hybrid gene, NPM-ALK, created by the t(2,5)(p23,q35) chromosomal translocation encodes part of the nucleolar phosphoprotein, nucleophosmin (NPM), joined to the entire cytoplasmic portion of the anaplastic lymphoma kinase (ALK) receptor tyrosine kinase. As a consequence, the ALK gene comes under the control of the NPM promoter, which induces a permanent and ubiquitous transcription of the NPM-ALK hybrid gene, resulting in the production of a 80 kDa NPM-ALK chimeric protein. This translocation is found in anaplastic large cell lymphomas (ALCL). Reportedly, expression of ALK indicates a better prognosis. Approximately 5 % -10 % of non-small cell lung carcinomas also express ALK protein producing a cytoplasmic staining pattern. This MAb also reacts with blood vessels that serves as an internal positive control.

**Pathways:** [RTK Signaling](#)

## Application Details

---

**Application Notes:** The stated application concentrations are suggested starting amounts. Titration of the ALK antibody may be required due to differences in protocols and secondary/substrate sensitivity.\.  
IHC (FFPE): 1-2 µg/mL for 30 min at RT

**Restrictions:** For Research Use only

## Handling

---

**Buffer:** 1 mg/mL in 1X PBS, BSA free, sodium azide free

**Preservative:** Azide free

**Storage:** 4 °C, -20 °C

**Storage Comment:** Store the recombinant ALK antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).