

Datasheet for ABIN6655122
anti-TERT antibody (C-Term)



[Go to Product page](#)

6 Images

5 Publications

Overview

Quantity:	25 µL
Target:	TERT
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TERT antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Fluorescence Microscopy (FM)

Product Details

Purpose:	hTERT Antibody
Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a region near the carboxy terminal end of hTERT (accession number AF018167).
Isotype:	IgG
Cross-Reactivity (Details):	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum.
Purification:	Affinity Purified Anti-hTERT Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using synthetic peptide coupled to agarose beads.
Sterility:	Sterile filtered

Target Details

Target:	TERT
Alternative Name:	TERT (TERT Products)
Background:	<p>Synonyms: rabbit anti-TERT antibody, rabbit anti-Telomerase catalytic subunit antibody, hTERT, Telomerase reverse transcriptase, HEST2, Telomerase-associated protein 2, TP2, EST2, TCS1, TRT</p> <p>Background: Telomerase is a reverse transcriptase that adds telomeric repeats (TTAGGG)_n to chromosomal ends, compensating for the telomere shortening that occurs with DNA replication. In normal human somatic cells, telomerase is repressed and telomeres progressively shorten, leading to limited lifespan and senescence. Reactivation of telomerase activity is associated with human cancer and cell immortalization. Approximately 85 % of human cancers, including breast, prostate, stomach, bladder, colon, and liver cancer, have telomerase activity, whereas most normal somatic cells do not. The specificity of telomerase to human cancer has led to investigations of telomerase activity and expression as a tumor marker. For example, the presence of telomerase activity in human urine has been identified as a marker for human bladder carcinoma. Human telomerase consists of three major subunits: a catalytic protein subunit called hTERT (for human Telomerase Reverse Transcriptase), a template RNA called hTR, and telomerase-associated protein (TEP-1). TERT and hTR are minimally required to reconstitute telomerase activity in vitro. In human cells, hTR is constitutively expressed. TERT transcription is a primary mechanism for regulation of telomerase activity.</p> <p>Gene Name: TERT</p>
Gene ID:	7015
NCBI Accession:	NP_001180305
UniProt:	O14746
Pathways:	Telomere Maintenance

Application Details

Application Notes:	Immunoprecipitation_Dilution: IP 2µL per mg lysate ELISA_Dilution: 1:10,000 - 1:50,000 Immunohistochemistry_Dilution: 1:500 IF_Microscopy_Dilution: 1:500 Western_Blots_Dilution: 1:500
Comment:	Suggested Applications: ChIP, FISH, IHC, Multiplex

Application Details

Anti-Telomerase catalytic subunit antibody has been tested for use in ELISA, immunoblotting, immunoprecipitation, and immunofluorescence microscopy. In these assays, the antibody detects ectopically-expressed hTERT and high levels of endogenous hTERT. A SY5Y cell nuclear extract can be used as a positive control. This antibody primarily detects hTERT, but several non-specific bands appear on immunoblots. In immunofluorescence microscopy assays, staining with anti-TERT-16 was specific to the nuclei of cells with ectopic TERT expression. In immunoblot assays, whole cell or nuclear extracts were loaded at a concentration of 100 µg protein per well. A working dilution of 1:500 anti-TERT antibody was used followed by a 1:3,000 dilution of HRP goat anti-rabbit IgG as the secondary antibody. For immunofluorescence microscopy staining, a working dilution of 1:500 was used followed by a 1:200 dilution of rhodamine-conjugated donkey anti-rabbit IgG as a secondary antibody. Immunoprecipitation was performed using 20 µL of protein A beads and 2 µL of the anti-TERT serum per 1 mg protein from cell lysate. A working dilution of 1:500 is also suggested for immunohistochemistry. To detect TERT, fix cells in 2% paraformaldehyde (in PBS) for 10'. Wash the slides twice in PBS for 5' each. Permeabilize the cells in 0.5% NP-40 for 10'. Wash as before in PBS. Block the cells using PBG buffer (0.2% cold water fish gelatin (Sigma G-7765) and 0.5% BSA in PBS) for 20' at room temperature. Incubate in primary antibody (diluted in PBG) for 1-2 hours at RT or overnight at 4°C. Wash the slides three times in PBG for 5' each. Incubate with secondary antibody (diluted in PBG) for 1 hour at RT in the dark. Wash the slides three times in PBG for 5' each. Mount in DAPI-containing medium.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Stabilizer: None

Preservative: 0.01 % (w/v) Sodium Azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated

Handling

above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

Expiry Date: 3 months

Publications


Product cited in: Bernal, Zafon, Domínguez, Bertran, Tusell: "Generation of Immortalised But Unstable Cells after hTERT Introduction in Telomere-Compromised and p53-Deficient vHMECs." in: **International journal of molecular sciences**, Vol. 19, Issue 7, (2018) ([PubMed](#)).

Eitan, Braverman, Tichon, Gitler, Hutchison, Mattson, Priel: "Excitotoxic and Radiation Stress Increase TERT Levels in the Mitochondria and Cytosol of Cerebellar Purkinje Neurons." in: **Cerebellum (London, England)**, Vol. 15, Issue 4, pp. 509-17, (2017) ([PubMed](#)).

Radan, Hughes, Teichroeb, Vieira Zamora, Jewer, Postovit, Betts: "Microenvironmental regulation of telomerase isoforms in human embryonic stem cells." in: **Stem cells and development**, Vol. 23, Issue 17, pp. 2046-66, (2014) ([PubMed](#)).


Theurillat, Udeshi, Errington, Svinkina, Baca, Pop, Wild, Blattner, Groner, Rubin, Moch, Privé, Carr, Garraway: "Prostate cancer. Ubiquitylome analysis identifies dysregulation of effector substrates in SPOP-mutant prostate cancer." in: **Science (New York, N.Y.)**, Vol. 346, Issue 6205, pp. 85-9, (2014) ([PubMed](#)).

Wu, Dudognon, Nguyen, Hillion, Pendino, Tarkanyi, Aradi, Lanotte, Tong, Chen, Ségal-Bendirdjian : "Immunodetection of human telomerase reverse-transcriptase (hTERT) re-appraised: nucleolin and telomerase cross paths." in: **Journal of cell science**, Vol. 119, Issue Pt 13, pp. 2797-806, (2006) ([PubMed](#)).




lot 21422, 1:2000 overnight

lot 25694, 1:2000 overnight



untreated

500 mM H2O2



Immunofluorescence

Image 1. anti-hTERT antibody-Immunofluorescence. Two different lots of anti-hTERT antibody were used to stain hTERT on hTERT-over-expressing fibroblasts. Cells were fixed in 4% PFA (in PBS) for 10 min and frozen in -80 after 3 min air-drying before incubation with anti hTERT 1:2000 overnight and staining with a 1:2000 dilution of Alexafluor488 secondary Ab. Confocal images provided by G. Saretzki, Institute for Ageing and Health, Newcastle University, UK. See Ahmed et. Al. for more information.

Immunofluorescence

Image 2. Anti-hTERT Antibody - Immunofluorescence Microscopy anti hTERT antibody-Immunofluorescence# anti hTERT antibody was used to stain hTERT on hTERT-over-expressing fibroblasts. Cells were untreated (Left) or treated (Right) with 500 uM H2O2, fixed in 4% PFA (in PBS) for 10 min and frozen in -80 after 3 min air-drying before staining with anti hTERT 1:2000 overnight. Confocal images provided by G. Saretzki, Institute for Ageing and Health, Newcastle University, UK. See Ahmed et. Al. for more information.

Western Blotting

Image 3. Rabbit anti-hTERT WB Western Blot of Rabbit anti-Telomerase catalytic subunit antibody. Lane 1: HeLa HV. Lane 2: HeLa 6-2. Load: 10 µg per lane. Primary antibody: Telomerase catalytic subunit antibody at 1:1,000 for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:40,000 for 45 min at RT. Block: Blocking Buffer for Fluorescent Western Blotting . Predicted/Observed size: 127 kDa, 127 kDa for Telomerase catalytic subunit.

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN6655122.