

# Datasheet for ABIN6655531 anti-CD4 antibody (FITC)





#### Overview

Quantity:	500 μL
Target:	CD4
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD4 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS), Fluorescence Microscopy (FM)

#### **Product Details**

Purpose:	CD4 Fluorescein Antibody
Immunogen:	Anti-CD4 Antibody (Monoclonal) was produced by repeated immunizations with PHA-stimulated human peripheral blood mononuclear cells (PBMC).
Clone:	RPA-T4
Isotype:	IgG1 kappa
Cross-Reactivity (Details):	Reactivity is observed against human CD4 and chimpanzee.
Purification:	Fluorescein conjugated CD4 Monoclonal Antibody was Protein G Purified from tissue culture supernatant and is directed against human CD4.
Sterility:	Sterile filtered
Labeling Ratio:	2-8


## **Target Details**


Target:	CD4
Alternative Name:	CD4 (CD4 Products)
Background:	Synonyms: T-cell surface glycoprotein CD4, T-cell surface antigen T4/Leu-3, CD4, FITC anti-
	human CD4, RPA-T4, FITC, T4
	Background: CD4 is an Immunoglobulin Superfamily protein and an accessory protein for MHC
	class-II antigen/ T-cell receptor interaction. It may regulate T-cell activation. It plays a more
	general role in mediating cell recognition events than merely those of cellular immune
	response. The T4 Molecule serves as a receptor for the human immunodeficiency virus. A Type
	I membrane protein, it is expressed on T lymphocytes, B-cells, macrophages, and granulocytes
	It is also expressed in a developmentally regulated manner in specific regions of the brain.
	Gene Name: CD4
Gene ID:	920
NCBI Accession:	NP_000607
UniProt:	P01730
Pathways:	TCR Signaling, Maintenance of Protein Location, CXCR4-mediated Signaling Events
Application Details	
Application Notes:	Flow_Cytometry_Dilution: 10 μL/10 <sup>6</sup> cells (0.1 μg)
	IF_Microscopy_Dilution: User Optimized
Comment:	Anti-CD4 has been tested in Flow Cytometry (Cell Surface) and is useful for
	Immunofluorescence. Researchers should determine optimal titers for applications that are no
	stated.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Buffer: 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
	Stabilizer: 0.1 % Gelatin
	Preservative: 0.09 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

#### Handling

	should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. DO NOT FREEZE. This product is light sensitive.
Expiry Date:	6 months


#### **Images**






#### **Flow Cytometry**

Image 1. Flow Cytometry - Mouse anti-HUMAN CD3 FITC Flow Cytometry of Mouse anti-HUMAN CD3 antibody Fluorescein conjugated. Cells: 10^6 Human peripheral blood lymphocytes. Stimulation: none. Antibody: (BLUE) Fluorescein Anti-CD3 mouse secondary antibody, (ORANGE) Fluorescein Anti-CD3 mouse control antibody.



### **Flow Cytometry**

**Image 2.** Flow Cytometry - Mouse anti-HUMAN CD3 FITC Flow Cytometry of Mouse anti-HUMAN CD3 antibody Fluorescein conjugated. Cells: 10^6 Human peripheral blood lymphocytes. Stimulation: none. Antibody: (Black) FITC Mouse isotype control; (Green) Fluorescein Anti-CD3 mouse secondary antibody using 5 ul.



#### **Flow Cytometry**

Image 3. Flow Cytometry - Mouse anti-HUMAN CD4 FITC Flow Cytometry of Mouse anti-HUMAN CD4 antibody Fluorescein conjugated. Cells: 10^6 Human peripheral blood lymphocytes. Stimulation: none. Antibody: (GRAY) 1µg FITC Mouse isotype control; (BLUE) Fluorescein Anti-CD3 mouse secondary antibody using 5 ul (1µg).

Please check the product details page for more images. Overall 4 images are available for ABIN6655531.