

## Datasheet for ABIN6656075

# anti-HMOX1 antibody





()	ve	rvi	6	W
$\sim$	v C	1 V I	$\sim$	v v

Quantity:	100 μg
Target:	HMOX1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMOX1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Fluorescence Microscopy (FM)
Product Details	
Purpose:	HO-1 Antibody
Immunogen:	HO-1 Antibody was produced from whole rabbit serum prepared by repeated immunizations
	raised against a human heme-oxygenase (HO-1) synthetic multiple antigenic synthetic peptide.
Isotype:	IgG
Cross-Reactivity (Details):	A BLAST analysis was used to suggest cross-reactivity with HO-1 from Human, Mouse, Rat and
	Dog based on 100 % homology with the immunizing sequence.
Purification:	Anti-HO-1 Antibody was purified by affinity chromatography.
Sterility:	Sterile filtered
Target Details	
Target:	HMOX1

### **Target Details**

Alternative Name:	HO-1 (HMOX1 Products)
Background:	Synonyms: Hsp32, HMOX1, Heme oxygenase 1, H0, H01  Background: Heme-oxygenase is a ubiquitous enzyme that catalyzes Heme-oxygenase is a ubiquitous enzyme that catalyzes.  Gene Name: HMOX1
Gene ID:	3162
NCBI Accession:	NP_002124
UniProt:	P09601
Pathways:	Transition Metal Ion Homeostasis, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, SARS-CoV-2 Protein Interactome

## Application Details

Application Notes:	Immunohistochemistry_Dilution: User Optimized  IF_Microscopy_Dilution: 1:100  Western_Blot_Dilution: 1:1000  Other: ICC: 1:100
Comment:	Anti-HO-1 Antibody has been tested in WB, IHC, IF. Expect a band approximately ~32kDa protein, corresponding to the molecular mass of HO-1 on SDS Page immunoblots. Specific conditions for reactivity should be optimized by the end user.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Format: Buffer:	Liquid  Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  Stabilizer: 50 % (v/v) Glycerol  Preservative: 0.09 % (w/v) Sodium Azide
	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 50 % (v/v) Glycerol
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 50 % (v/v) Glycerol Preservative: 0.09 % (w/v) Sodium Azide


### Handling

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiry Date:

12 months

#### **Images**



#### **Western Blotting**

**Image 1.** HO-1 Western Blot. Western Blot of rabbit anti-HO-1 antibody. Lane 1: Human cell line mix. Lane 2: none. Load: 35 μg per lane. Primary antibody: HO-1 antibody at 1:1000 for overnight at 4°C. Secondary antibody: rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 32.8 kDa, ~35 kDa for HO-1. Other band(s): none.