

Datasheet for ABIN6657494

anti-ATM antibody (pSer1981)

5 Images

118 Publications

[Go to Product page](#)

Overview

Quantity:	25 µL
Target:	ATM
Binding Specificity:	AA 1974-1988, pSer1981
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This ATM antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunoprecipitation (IP), Immunofluorescence (IF), Flow Cytometry (FACS), Fluorescence Microscopy (FM)

Product Details

Purpose:	ATM phospho S1981 Antibody
Immunogen:	Anti-ATM phospho S1981 Antibody was produced from a synthetic peptide S-L-A-F-E-E-G-Sp-Q-S-T-T-I-S-S corresponding to aa 1974-1988 of human ATM.
Clone:	10H11-E12
Isotype:	IgG1 kappa
Cross-Reactivity (Details):	This monoclonal anti-ATM antibody recognizes the phosphorylated epitope in native and over-expressed proteins found in various tissues and extracts.
Purification:	Anti-ATM phospho S1981 Monoclonal Antibody is directed against human ATM and is useful in determining its presence in various assays.

Product Details

Sterility: Sterile filtered

Target Details

Target: ATM

Alternative Name: ATM ([ATM Products](#))

Background: Synonyms: mouse anti-ATM antibody, mouse anti-ATMpS1981 antibody, mouse anti- ATM pS1981 antibody, DKFZp781A0353 antibody, Human phosphatidylinositol 3 kinase homolog antibody, MGC74674 antibody, Serine protein kinase ATM antibody, T cell prolymphocytic leukemia antibody

Background: Anti ATM pS1981 Antibody is a phospho site specific antibody and recognizes the product of the ATM gene that is mutated in the hereditary disease ataxia-telangiectasia. ATM codes for a protein kinase that acts as a master regulator of cellular responses to DNA double-strand breaks. ATM is normally inactive and the question of how it is activated in the event of DNA damage (due to ionizing radiation for instance) is central to understanding its function. ATM protein is now shown to be present in undamaged cells as an inactive dimer. Low doses of ionizing radiation, which induce only a few DNA breaks, activate at least half of the total ATM protein present, possibly in response to changes in chromatin structure. The ATM gene encodes a 370- kDa protein that belongs to the phosphoinositide 3-kinase (PI(3)K) superfamily, but which phosphorylates proteins rather than lipids. The 350-amino-acid kinase domain at the carboxy terminus of this large protein is the only segment of ATM with an assigned function. Exposure of cells to IR triggers ATM kinase activity, and this function is required for arrests in G1, S and G2 phases of the cell cycle. Several substrates of the ATM kinase participate in these IR-induced cell-cycle arrests. These include p53, Mdm2 and Chk2 in the G1 checkpoint, Nbs1, Brca1, FancD2 and SMC1 in the transient IR-induced S-phase arrest, and Brca1 and hRad17 in the G2/M checkpoint. Ideal for Cancer, Cell Signaling, Chromatin, Neuroscience and Signal Transduction research.

Gene Name: ATM

Gene ID: 472

NCBI Accession: [NP_000042](#)

UniProt: [Q13315](#)

Pathways: [p53 Signaling](#), [Apoptosis](#), [DNA Damage Repair](#), [Inositol Metabolic Process](#), [Positive Regulation of Response to DNA Damage Stimulus](#)

Application Details

Application Notes: Immunoprecipitation_Dilution: User Optimized
ELISA_Dilution: 1:20,000 - 1:100,000
Immunohistochemistry_Dilution: Not Recommended
Flow_Cytometry_Dilution: 5 µg/mL
IF_Microscopy_Dilution: 1:100 - 1:500
Western_Blot_Dilution: 1:200 - 1:2,000
Other: User Optimized

Comment: Suggested Applications:
Suggested_Applications: Biochemical Assay, ChIP, FISH, IHC, IP, Multiplex
Other_Performance_Data: iFISH at 7.5µg/ml

Protein A Purified Mab anti-ATM has been tested by ELISA, FC, IF, and Western blotting against both the native and recombinant forms of the protein. The antibody immunoprecipitates ATM from irradiated human and transfected mouse cells. By immunofluorescence, foci are detected in irradiated human and mouse fibroblasts. This antibody is not recommended for immunohistochemistry. Instead, for IHC, use the clone 7C10D8 (item# 200-301-500).

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer: None
Preservative: 0.01 % (w/v) Sodium Azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

Handling

Expiry Date: 12 months

Publications

Product cited in: Pan, Wu, Yeh: "ATM Inhibitor Suppresses Gemcitabine-Resistant BTC Growth in a Polymerase θ Deficiency-Dependent Manner." in: **Biomolecules**, Vol. 10, Issue 11, (2024) ([PubMed](#)).


Li, Gai, Li, Yang, Yu: "DNA-PK participates in pre-rRNA biogenesis independent of DNA double-strand break repair." in: **Nucleic acids research**, Vol. 52, Issue 11, pp. 6360-6375, (2024) ([PubMed](#)).

Xin, Gai, Ma, Li, Li, Yu: "Pre-rRNA Facilitates TopBP1-Mediated DNA Double-Strand Break Response." in: **Advanced science (Weinheim, Baden-Wuerttemberg, Germany)**, Vol. 10, Issue 28, pp. e2206931, (2023) ([PubMed](#)).

Chen, Li, Iemura, Tanaka: "Oxidative stress induces chromosomal instability through replication stress in fibroblasts from aged mice." in: **Journal of cell science**, Vol. 136, Issue 11, (2023) ([PubMed](#)).


Gritti, Basso, Rinchai, Corigliano, Pivetti, Gaviraghi, Rosano, Mazza, Barozzi, Roncador, Parmigiani, Legube, Parazzoli, Cittaro, Bedognetti, Mondino, Segalla, Tonon: "Loss of ribonuclease DIS3 hampers genome integrity in myeloma by disrupting DNA:RNA hybrid metabolism." in: **The EMBO journal**, Vol. 41, Issue 22, pp. e108040, (2022) ([PubMed](#)).

There are more publications referencing this product on: [Product page](#)




Immunofluorescence

Image 1. Anti-ATM Monoclonal Antibody - Immunofluorescence Microscopy anti-ATM pS1981 mouse monoclonal antibody (Catalog # 200-301-400) detects ATM phosphorylated on Ser 1981 by Indirect immunofluorescence microscopy. Shown are hTCEpi cells (courtesy of Dr. Danielle Robertson) infected with HSV-1 at MOI 5.0 and fixed at 8 hpi with 3% paraformaldehyde/2% sucrose for 10 min. After rinsing, cells were permeabilized with 0.5% Triton X-100 for 5 min, blocked with 3% BSA for 30 min, and stained with primary anti-ATM pS1981 antibody overnight at 5 µg/mL (1:200). Secondary staining was performed with Alexa Fluor 594 anti-mouse antibody. Images were taken with Olympus AX70 compound epifluorescence microscope equipped with Spot RT Slider camera. Experiment was performed by Oleg Alekseev in the laboratory of Dr. Jane Azizkhan-Clifford at Drexel University College of Medicine.



Immunofluorescence

Image 2. Anti-ATM Monoclonal Antibody - Immunofluorescence Microscopy Anti ATM antibody showing overlay of anti-ATM pS1981 staining. Cells were fixed 15 min after 5 Gy (IR+) of irradiation, then labeled with antibody. See Kitagawa et al. for additional details.



Western Blotting

Image 3. Anti-ATM Monoclonal Antibody - Western Blot Anti ATM Mab with human derived HEK293 cells treated with doxorubicin using Protein A Purified Mab anti-ATM Protein Kinase pS1981(clone 10H11.E12). A 370 kDa band corresponding to phosphorylated ATM is detected (arrowhead, lane 2). The lysate was prepared with HALT phosphatase inhibitor (Pierce). Pre-incubation of peptide with 50 µg of immunizing phospho peptide negates specific staining (lane 1). Approximately 30 µg of lysate was added to each lane of an SDS-PAGE gel under non-reducing conditions. The protein was transferred to nitrocellulose using standard methods. After blocking the membrane was probed with the primary antibody diluted 1:500 overnight at 4°C followed by washes and reaction with a 1:10,000 dilution of 800 conjugated Gt-a-Mouse IgG [H&L] (code 610-132-121) for 40 min at room temperature. LICOR's Infrared Imaging System was used to scan and process the image. Other detection systems will yield similar results.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN6657494.