

Datasheet for ABIN6657569
anti-Hemagglutinin antibody



[Go to Product page](#)

1 Image 5 Publications

Overview

Quantity:	100 µg
Target:	Hemagglutinin (HA)
Reactivity:	Influenza A Virus H5N1
Virus Strain:	A/Vietnam/1203/04
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Hemagglutinin antibody is un-conjugated
Application:	ELISA, Immunoprecipitation (IP), Western Blotting (WB), Neutralization (Neut), Immunohistochemistry (IHC)

Product Details

Purpose:	H5N1 Antibody VN04-13
Immunogen:	Hemagglutinin of A/Vietnam/1203/04 Influenza Virus (VN04-13) monoclonal antibody was produced by intraperitoneal immunization of BALB/c mice with concentrated purified virus preparation containing hemagglutinin (HA) protein of influenza A virus [strain A/Vietnam/1203/04 (H5N1)] using the modification of the method described by Kohler and Milstein. Each mouse received two immunizations of 15 mug HA with incomplete Freund's adjuvant, administered 3 week apart.
Clone:	14C5
Isotype:	IgG2a kappa
Cross-Reactivity (Details):	VN04-13 monoclonal antibody did not cross-react with influenza viruses of other HA subtypes.

Product Details

Purification:	H5N1 Antibody was purified from tissue culture supernatant fluid by Protein A chromatography and is specific for H5 hemagglutinin (HA) protein of influenza A virus [strain A/Vietnam/1203/04 (H5N1)].
Sterility:	Sterile filtered

Target Details

Target:	Hemagglutinin (HA)
Alternative Name:	Hemagglutinin (HA Products)
Target Type:	Influenza Protein
Background:	Synonyms: mouse anti-H5N1 Antibody, mouse anti-VN04-13 Antibody, H5HA antibody, Hemagglutinin 5 antibody, H5N1 antibody Background: Hemagglutinin of A/Vietnam/1203/04 Influenza Virus (VN04-13) Antibody raised against the hemagglutinin (HA) surface glycoprotein of the A/Vietnam/1203/04 (H5N1) influenza virus. Generally referred to as "bird flu", the H5N1 influenza A virus has been documented in poultry and humans across ten Eurasian countries, from Japan in the north to Indonesia in the south. Without immunity, humans would have no protection against H5N1 influenza viruses, which could potentially cause a catastrophic pandemic influenza. This antibody, directed against the HA surface glycoprotein of the A/Vietnam/1203/04 (H5N1) influenza virus, is intended to further our understanding of the mechanisms underlying antigenic variation and evolution of novel variants. The major functions of HA include receptor-binding and fusion activities, but there may also be a structural role for HA in viral particle formation. Following attachment of HA to surface receptors on susceptible cells, the influenza virus enters the cell via endocytosis and membrane fusion. Gene Name: HA
Gene ID:	159144921
UniProt:	A8UDQ2

Application Details

Application Notes:	Neutralization_Dilution: User Optimized
	Immunoprecipitation_Dilution: User Optimized
	ELISA_Dilution: 1:5,000
	Immunohistochemistry_Dilution: User Optimized
	Western_Blot_Dilution: User Optimized

Application Details

Comment: Hemagglutinin of A/Vietnam/1203/04 Influenza Virus (VN04-13) monoclonal antibody can be used for hemagglutination inhibition (HI) assays to provide antigenic characterization of the influenza A viruses of the H5 HA subtype. This monoclonal antibody is suitable for virus neutralization assays (in cell culture and in embryonated chicken eggs), ELISA, immunoprecipitation, immunohistochemistry and western blotting.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Stabilizer: None

Preservative: 0.01 % (w/v) Sodium Azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiry Date: 12 months

Publications

Product cited in: Reed, Yen, DuBois, Bridges, Salomon, Webster, Russell: "Amino acid residues in the fusion peptide pocket regulate the pH of activation of the H5N1 influenza virus hemagglutinin protein." in: **Journal of virology**, Vol. 83, Issue 8, pp. 3568-80, (2009) ([PubMed](#)).


Kaverin, Rudneva, Govorkova, Timofeeva, Shilov, Kochergin-Nikitsky, Krylov, Webster: "Epitope mapping of the hemagglutinin molecule of a highly pathogenic H5N1 influenza virus by using monoclonal antibodies." in: **Journal of virology**, Vol. 81, Issue 23, pp. 12911-7, (2007) ([PubMed](#)).

Govorkova, Webby, Humberd, Seiler, Webster: "Immunization with reverse-genetics-produced H5N1 influenza vaccine protects ferrets against homologous and heterologous challenge." in: **The Journal of infectious diseases**, Vol. 194, Issue 2, pp. 159-67, (2006) ([PubMed](#)).

Hanson, Boon, Lim, Webb, Ooi, Webby: "Passive immunoprophylaxis and therapy with humanized monoclonal antibody specific for influenza A H5 hemagglutinin in mice." in: **Respiratory research**, Vol. 7, pp. 126, (2006) ([PubMed](#)).

Hoffmann, Lipatov, Webby, Govorkova, Webster: "Role of specific hemagglutinin amino acids in the immunogenicity and protection of H5N1 influenza virus vaccines." in: **Proceedings of the National Academy of Sciences of the United States of America**, Vol. 102, Issue 36, pp. 12915-20, (2005) ([PubMed](#)).

Images



Cross reactivity of anti-A/Vietnam/1203/2004 (H5N1) HA monoclonal antibodies with H5N1 influenza viruses in HI assay.							
HA Clade	H5N1 Influenza Virus	HI titers with anti-HA monoclonal antibodies:					
HS Ref.		VN04-2	VN04-8	VN04-9	VN04-10	VN04-13	VN04-16
North American	A/Tenn/South Africa/61	100	<	<	<	<	<
	A/Chicken/Pennsylvania/1370/83	3200	<	25600	200	3200	<
	A/Mallard/Pennsylvania/10218/84	800	<	200	6400	25600	400
Clade 0	A/Chicken/Hidalgo/28159-233/94	<	<	200	100	1600	<
	A/Mallard/Arkansas/1/2001	1600	<	200	400	3200	100
	A/Hong Kong/156/97	6400	<	25600	6400	25600	400
Clade 1	A/Hong Kong/481/97	6400	<	1600	1600	12800	100
	A/Duck/Singapore/3/97	200	<	200	800	6400	200
	A/Goose/Hong Kong/437-4/99	6400	<	6400	1600	6400	200
Clade 2.1.2	A/Vietnam/1194/2004	3200	1600	12800	3200	6400	1600
	A/Vietnam/1203/2004	6400	1600	12800	3200	6400	1600
	A/Vietnam/HN30409/2005	6400	3200	3200	3200	6400	1600
Clade 2.1.3	A/Hong Kong/213/2003	6400	3200	400	3200	800	3200
	A/Indonesia/9/2005	3200	<	800	25600	200	6400
	A/Indonesia/PA03/2003	800	3200	200	3200	1600	1600
Clade 2.2	A/Duck/HUNVIR/504/2004	1600	<	3200	1600	<	400
	A/Duck/HK/1304/2004	<	1600	<	3200	1600	1600
	A/Chicken/Los Angeles/BBV/ET/02/2004	100	<	100	3200	3200	400
Clade 2.3.4	A/Chicken/Malang/BBV/ET/VI/2004	3200	3200	<	3200	3200	1600
	A/Whooper Swan/Mongolia/244/2005	<	1600	<	3200	1600	1600
	A/Turkey/15/2006	100	<	<	3200	<	400
Clade 2.4	A/Bar-headed goose/Qinghai/1A/2005	100	6400	<	6400	12800	3200
	A/Duck/Hunan/15/2004	1600	<	3200	1600	<	400
	A/Duck/Laos/3295/2006	<	<	400	1600	100	100
Clade 2.4	A/Chicken/Malaysia/95/2006	100	<	400	800	100	100
	A/Common mague/Hong Kong/845/2006	<	<	200	400	<	100
Clade 2.4	A/Duck/Guangxi/13/2004	<	1600	<	3200	1600	1600

Hemagglutination-inhibition (HI) testing was performed with 0.5% chicken red blood cells by standard method. < = less than 1:100.