

# Datasheet for ABIN6658223

# anti-GABRA6 antibody (Cytoplasmic Loop)





Go to Product page

( )	VE	IN	/1	1	Α
1 )	<b>\/</b> ⊢	3 I /	/ 1 (	_,	//\
$\sim$	٧ ر	, I 1	/ I'	$\overline{}$	v v

Overview		
Quantity:	100 μL	
Target:	GABRA6	
Binding Specificity:	Cytoplasmic Loop	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GABRA6 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Purpose:	GABA(A) Receptor alpha 6 Antibody	
Immunogen:	Anti-GABA(A) Receptor alpha 6 Antibody was produced in rabbit by repeated immunizations	
	with recombinant fusion proteins from the cytoplasmic loop of the alpha 6 subunit.	
Isotype:	IgG	
Cross-Reactivity (Details):	Anti-GABA(A) Receptor alpha 6 Antibody is directed against rat GABA(A) Receptor alpha 6.	
Purification:	The antibody was affinity purified from monospecific antiserum by immunoaffinity purification.	
Target Details		
Target:	GABRA6	
Alternative Name:	GABA(A) Receptor alpha 6 (GABRA6 Products)	

#### Target Details

Background:

Synonyms: Gamma-aminobutyric acid receptor subunit alpha-6, GABA(A) receptor subunit alpha-6

Background: Anti-GABA(A) Receptor alpha 6 Antibody detects GABA(A) Receptor alpha 6. Gamma-aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system, causing a hyperpolarization of the membrane through the opening of a Cl-channel associated with the GABAA receptor (GABAA-R) subtype. GABAA-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and substance abuse. The GABAA-R is a multimeric subunit complex. To date six  $\alpha$ s, four  $\beta$ s and four  $\gamma$ s, plus alternative splicing variants of some of these subunits, have been identified. Injection in occytes or mammalian cell lines of cRNA coding for  $\alpha$ - and  $\beta$ -subunits results in the expression of functional GABAA-Rs sensitive to GABA. However, coexpression of a  $\gamma$ -subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different  $\alpha$ -subunits of the receptor. GABA(A) receptor alpha 6 antibody is ideal for investigators involved in Neuroscience.

Gene Name: GABRA6

Gene ID:

29708

NCBI Accession:

NP\_068613

UniProt:

P30191

Pathways:

Synaptic Membrane

#### **Application Details**

App	lication	Notes:

Optional[Neutralization\_Dilution]: 1:1000

Comment:

Anti-GABA(A) Receptor alpha 6 (Rabbit) antibody is tested for use in Western Blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band of approximately 57 kDa in size corresponding to the alpha 6 subunit of the GABA(A) receptor in the appropriate cell lysate or extract.

Restrictions:

For Research Use only

#### Handling

Format:

Liquid

Buffer:


Buffer: 0.01 M HEPES, 0.15 M Sodium Chloride, pH 7.5

#### Handling

	Stabilizer: 0.1 mg/mL Bovine Serum Albumin (BSA) - IgG and Protease free, 50 % (v/v) Glycerol
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at -20° C prior to opening. This product is stable at 4° C as an undiluted liquid. For extended storage, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Dilute only prior to immediate use.
Expiry Date:	12 months

## **Images**

## Anti-GABA<sub>A</sub> Receptor, α<sub>6</sub>-Subunit



Western blot of 5-7  $\mu g$  of mouse cerebellum lysates from wild type (control) and  $\alpha_{\rm g}$  knockout ( $\alpha_{\rm g}$  K/O) animals showing specific immunolabeling of the ~57k  $\alpha_{\rm g}$ -subunit of the GABA<sub>A</sub>-R in the wild type but not in the  $\alpha_{\rm g}$  K/O animals.

#### **Western Blotting**

Image 1. Western Blot of Anti-GABA(A) Receptor alpha 6 (Rabbit) Antibody - 600-401-D48 Western Blot of Rabbit anti-GABA(A) Receptor alpha 6 antibody. Lane 1: mouse forebrain lysates from Wild Type. Lane 2: mouse forebrain lysates from Wild Type  $\alpha$ 6-knockout ( $\alpha$ 6-K/O). Load: 10  $\mu$ g per lane. Primary antibody: GABAA-R antibody at 1:400 for overnight at 4°C. Secondary antibody: rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: ~57kDa/~56kDa for  $\alpha$ 6-subunit of the GABAA-R. Other band(s): none.