
antibodies .- online.com







anti-CRISPR-Cas9 (C-Term) antibody (DyLight 549)





\sim					
	VE	۲۱د	/1/	$\triangle V$	٨
\cup	V	ر ار ا	יו ע	U V	V


Quantity:	100 μg	
Target:	CRISPR-Cas9	
Binding Specificity:	C-Term	
Reactivity:	Streptococcus pyogenes	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	DyLight 549	
Application:	Western Blotting (WB), ELISA, Fluorescence Microscopy (FM)	
Product Details		
Immunogen:	Immunogen: Cas-9 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a region near the C-terminal of Streptococcus pyogenes Cas-9 protein. Immunogen Type: Peptide	
Isotype:	IgG	
Purification:	Anti-Cas9 antibody is directed against Streptococcus pyogenes Cas-9. The product was affinity purified from monospecific antiserum by immunoaffinity chromatography.	
Labeling Ratio:	4.8	
Target Details		
Target:	CRISPR-Cas9	

Target Details

Alternative Name:	Cas 9		
Background:	Synonyms: rabbit anti-Cas9 antibody DyLight™ 549 conjugation, DyLight™ 549 conjugated rabbi		
	anti-Cas9 antibody, Cas-9, Cas 9, CRISPR, CRISPR-associated endonuclease Cas9/Csn1,		
	SpCas9, SpyCas9, csn1, Cas-9 Antibody		
	Background: The Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and		
	CRISPR Associated (Cas) system was discovered in bacteria as a defense against foreign DNA,		
	even being believed to act as a sort of prokaryotic immune system. The Cas9 protein is		
	becoming a a useful tool in the field of genomic editing for its ability to induce site-directed		
	double strand breaks in DNA. It can cleave almost any sequence complementary to the guide		
	RNA and has recently been used for modifying the genome of human embryos for the first		
	time.		
	Gene Name: cas9		
Gene ID:	901176		
NCBI Accession:	NP_269215		
UniProt:	Q99ZW2		
Application Details			
	Application Natar Coop DulimetTME 40 Conjugated Aptihody book book tooted for you in ELICA		
Application Notes:	Application Note: Cas9 DyLight™549 Conjugated Antibody has been tested for use in ELISA,		
	Immunofluorescence, and western blotting. Specific conditions for reactivity should be		
	optimized by the end user. Expect a band ~158 kDa in size corresponding to Cas-9 protein by		
	western blotting in the appropriate cell lysate or extract.		
	ELISA Dilution: 1,25,000-1:30,000 Western Blot Dilution: 1 µg/mL		
	IF Microscopy Dilution: 1 µg/mL		
	Other Performance Data: User Optimized		
Restrictions:	For Research Use only		
Handling			
Format:	Lyophilized		
Reconstitution:	Reconstitution Volume: 100 μL		
	Reconstitution Buffer: Restore with deionized water (or equivalent)		
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2		


	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Images



Western Blotting

Image 1. Western Blot of Rabbit Anti-Cas9 Antibody - Western Blot of Rabbit Anti-Cas9 Antibody. Lane 1: 293T non transfected cell lysate. Lane 2: 293T Cas9 over expressed cell lysate. Load: 15µg per lane. Primary Antibody cas9 used at 1µg/mL using ABIN925618 overnight at 4°C. Secondary Antibody: goat anti-rabbit HRP at 1:40,000 for 30 min at room temp. Expect: 158kDa.



Immunofluorescence

Image 2. Immunofluorescence of Rabbit Anti-Cas9 Antibody. - Immunofluorescence of Rabbit Anti-Cas9 Antibody. Cells: HeLa cells transfected with GFP+Cas9 or GFP+CD-3. Fixation: Formaldehyde fixed, permeabilized Triton X-100. Primary: Anti-Cas9 used at 1µg/mL. Secondary: Donkey anti-Rabbit at 1µg/mL, staining seen in red. Bottom images show GFP stain overlay.