



Datasheet for ABIN6658290

## anti-CRISPR-Cas9 (C-Term) antibody (DyLight 800)



[Go to Product page](#)

### 2 Images

#### Overview

Quantity:	100 µg
Target:	CRISPR-Cas9
Binding Specificity:	C-Term
Reactivity:	Streptococcus pyogenes
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	DyLight 800
Application:	Western Blotting (WB), ELISA, Fluorescence Microscopy (FM)

#### Product Details

Immunogen:	Immunogen: Cas-9 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a region near the C-terminal of Streptococcus pyogenes Cas-9 protein. Immunogen Type: Peptide
Isotype:	IgG
Purification:	Anti-Cas9 antibody is directed against Streptococcus pyogenes Cas-9. The product was affinity purified from monospecific antiserum by immunoaffinity chromatography.
Labeling Ratio:	2.4

#### Target Details

Target:	CRISPR-Cas9
---------	-------------

## Target Details

---

Alternative Name:	Cas 9
Background:	<p>Synonyms: rabbit anti-Cas9 antibody DyLight™ 800 conjugation, DyLight™ 800 conjugated rabbit anti-Cas9 antibody, Cas-9, Cas 9, CRISPR, CRISPR-associated endonuclease Cas9/Csn1, SpCas9, SpyCas9, csn1, Cas-9 Antibody</p> <p>Background: The Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and CRISPR Associated (Cas) system was discovered in bacteria as a defense against foreign DNA, even being believed to act as a sort of prokaryotic immune system. The Cas9 protein is becoming a a useful tool in the field of genomic editing for its ability to induce site-directed double strand breaks in DNA. It can cleave almost any sequence complementary to the guide RNA and has recently been used for modifying the genome of human embryos for the first time.</p> <p>Gene Name: cas9</p>
Gene ID:	901176
NCBI Accession:	<a href="#">NP_269215</a>
UniProt:	<a href="#">Q99ZW2</a>

## Application Details

---

Application Notes:	<p>Application Note: Cas9 DyLight™800 Conjugated Antibody has been tested for use in ELISA, Immunofluorescence, and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band ~158 kDa in size corresponding to Cas-9 protein by western blotting in the appropriate cell lysate or extract.</p> <p>ELISA Dilution: 1:25,000-1:30,000</p> <p>Western Blot Dilution: 1 µg/mL</p> <p>IF Microscopy Dilution: 1 µg/mL</p> <p>Other Performance Data: User Optimized</p>
Restrictions:	For Research Use only

## Handling

---

Format:	Lyophilized
Reconstitution:	<p>Reconstitution Volume: 100 µL</p> <p>Reconstitution Buffer: Restore with deionized water (or equivalent)</p>
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

## Handling

Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free  
0.01 % (w/v) Sodium Azide


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: RT, 4 °C, -20 °C


Storage Comment: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

## Images



### Western Blotting

**Image 1.** Western Blot of Rabbit Anti-Cas9 Antibody - Western Blot of Rabbit Anti-Cas9 Antibody. Lane 1: 293T non transfected cell lysate. Lane 2: 293T Cas9 over expressed cell lysate. Load: 15µg per lane. Primary Antibody cas9 used at 1µg/mL using ABIN925618 overnight at 4°C. Secondary Antibody: goat anti-rabbit HRP at 1:40,000 for 30 min at room temp. Expect: 158kDa.



### Immunofluorescence

**Image 2.** Immunofluorescence of Rabbit Anti-Cas9 Antibody. - Immunofluorescence of Rabbit Anti-Cas9 Antibody. Cells: HeLa cells transfected with GFP+Cas9 or GFP+CD-3. Fixation: Formaldehyde fixed, permeabilized Triton X-100. Primary: Anti-Cas9 used at 1µg/mL. Secondary: Donkey anti-Rabbit at 1µg/mL, staining seen in red. Bottom images show GFP stain overlay.