antibodies -online.com







Overview

Quantity:	100 μL
Target:	PKA
Binding Specificity:	AA 17-100
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKA antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from bovine PKA
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.

Target Details

Target:	PKA
Alternative Name:	PKA alpha + beta (PKA Products)

Target Details

_	
Background:	Synonyms: cAMP-dependent protein kinase catalytic subunit beta, PKA C-beta, PRKACB
	Background: Mediates cAMP-dependent signaling triggered by receptor binding to GPCRs. PKA
	activation regulates diverse cellular processes such as cell proliferation, the cell cycle, and
	differentiation and regulation of microtubule dynamics, chromatin condensation and
	decondensation, nuclear envelope disassembly and reassembly, as well as regulation of
	intracellular transport mechanisms and ion flux. Regulates the abundance of
	compartmentalized pools of its regulatory subunits through phosphorylation of PJA2 which
	binds and ubiquitinates these subunits, leading to their subsequent proteolysis (By similarity).
Gene ID:	282323
UniProt:	P05131
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months