

Datasheet for ABIN6698900

Goat anti-Guinea Pig IgG Antibody (DyLight 405) - Preadsorbed



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	IgG
Reactivity:	Guinea Pig
Host:	Goat
Clonality:	Polyclonal
Conjugate:	DyLight 405
Application:	Western Blotting (WB), FLISA, Fluorescence Microscopy (FM)

Product Details

Purpose:	Guinea Pig IgG (H&L) Antibody DyLight™ 405 Conjugated Pre-Adsorbed
Immunogen:	Guinea Pig IgG whole molecule
Isotype:	IgG
Cross-Reactivity (Details):	Minimal crossreactivity against Bv Ch Gt Ham Hs Hu Ms Rb Rt & Sh Serum Proteins
Characteristics:	Goat Anti-Guinea Pig IgG DyLight 405™ Conjugation, Goat Anti Guinea Pig IgG DyLight 405™ conjugated, Anti-Guinea Pig IgG DyLight Antibody generated in goat detects guinea pig IgG.
Purification:	PreadSORption: Pre-Adsorbed
Labeling Ratio:	4.9

Target Details

Target:	IgG
---------	-----

Target Details

Abstract: [IgG Products](#)

Target Type: Antibody

Background: Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75 % of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Application Details

Application Notes: FLISA_Dilution: >1:20,000
IF_Microscopy_Dilution: >1:5,000
Western_Blot_Dilution: >1:10,000
Other: User Optimized

Comment: The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitution Volume: 100 µL
Reconstitution Buffer: Restore with deionized water (or equivalent)

Concentration: 1.0 mg/mL

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum

Handling

Albumin (BSA) - Immunoglobulin and Protease free, 0.01 % (w/v) Sodium Azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store conjugated secondary antibody at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Conjugated Secondary Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiry Date: 12 months
