

Datasheet for ABIN6699008

Rabbit anti-Mouse IgG Antibody (DyLight 549)

1 Publication

Abstract:



Go to Product page

Overview	
Quantity:	100 μg
Target:	IgG
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	DyLight 549
Application:	Western Blotting (WB), FLISA, Fluorescence Microscopy (FM)
Product Details	
Purpose:	Mouse IgG (H&L) Antibody DyLight™ 549 Conjugated
Immunogen:	Mouse IgG, whole molecule
Isotype:	IgG
Characteristics:	rabbit anti-Mouse IgG Antibody DyLight™ 549 conjugation, rabbit anti-Mouse IgG DyLight™ 549 conjugated Antibody,Anti-Mouse IgG DyLight 549 Antibody generated in rabbit detects reactivity to Mouse IgG.
Labeling Ratio:	3.1
Target Details	
Target:	IgG

IgG Products

Target Details

Target Type:	Antibody
Background:	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G
	constitutes 75 $\%$ of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as
	well as fungi and facilitates their destruction or neutralization via agglutination (and thereby
	immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis.
	The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc recepto
	proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy
	and Light chains of the antibody molecule are present. Secondary Antibodies are available in a
	variety of formats and conjugate types. When choosing a secondary antibody product,
	consideration must be given to species and immunoglobulin specificity, conjugate type,
	fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
Application Details	
Application Notes:	FLISA_Dilution: >1:20,000
	IF_Microscopy_Dilution: >1:5,000
	Western_Blot_Dilution: >1:10,000
	Other: User Optimized
Comment:	This product is designed for immunofluorescence microscopy, fluorescence based plate
	assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex
	analysis, including multicolor imaging, utilizing various commercial platforms. The emission
	spectra for this DyLight™ conjugate match the principle output wavelengths of most common
	fluorescence instrumentation.
	Suggested Applications: IF
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 100 μL
	Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum
	Albumin (BSA) - Immunoglobulin and Protease free, 0.01 % (w/v) Sodium Azide

Handling

Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store conjugated secondary antibody at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Conjugated Secondary Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months
Publications	
Product cited in:	Yang, Ye, Dai, Li, Zhang, Xue, Zhu, Feng, Qin, Wang, Lei, Liao, Hao: "Phase separation of Epstein-Barr virus EBNA2 protein reorganizes chromatin topology for epigenetic regulation." in: Communications biology, Vol. 4, Issue 1, pp. 967, (2021) (PubMed).