

Datasheet for ABIN6699016

Donkey anti-Mouse IgG Antibody (DyLight 549) - Preadsorbed



[Go to Product page](#)

1 Image

1 Publication

Overview

Quantity:	100 µg
Target:	IgG
Reactivity:	Mouse
Host:	Donkey
Clonality:	Polyclonal
Conjugate:	DyLight 549
Application:	Western Blotting (WB), FLISA, Fluorescence Microscopy (FM)

Product Details

Purpose:	Mouse IgG (H&L) Antibody DyLight™ 549 Conjugated Pre-Adsorbed
Immunogen:	Mouse IgG whole molecule
Isotype:	IgG
Cross-Reactivity (Details):	Minimal crossreactivity against Bv Ch Gt GP Ham Hs Hu Rb Rt & Sh Serum Proteins
Characteristics:	Donkey anti-Mouse IgG DyLight 549™ Conjugated Antibody, Donkey anti Mouse IgG Antibody DyLight 549™ Conjugation, Anti-Mouse IgG DyLight549 Antibody generated in donkey detects reactivity to Mouse IgG.
Purification:	Preadsorption: Pre-Adsorbed
Labeling Ratio:	3.8

Target Details

Target: IgG

Abstract: [IgG Products](#)

Target Type: Antibody

Background: Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75 % of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Application Details

Application Notes: FLISA_Dilution: >1:20,000
IF_Microscopy_Dilution: >1:5,000
Western_Blot_Dilution: >1:10,000
Other: User Optimized

Comment: Anti-Mouse IgG DyLight549 Antibody has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation.
Suggested Applications: IF, IHC, Multiplex

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitution Volume: 100 µL
Reconstitution Buffer: Restore with deionized water (or equivalent)


Handling

Concentration:	1.0 mg/mL
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free, 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store conjugated secondary antibody at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Conjugated Secondary Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months

Publications

Product cited in: Scoles, Pflieger, Thai, Hansen, Dansithong, Pulst: "ETS1 regulates the expression of ATXN2." in: **Human molecular genetics**, Vol. 21, Issue 23, pp. 5048-65, (2013) ([PubMed](#)).

Images



Western Blotting

Image 1. Western Blot of 549 conjugated Donkey Anti-Mouse IgG Pre-Adsorbed secondary antibody. Lane 1: Mouse IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: 549 donkey secondary antibody at 1:1,000 for 60 min at RT. Blocking: ABIN925618 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Mouse IgG. Other band(s): None.