

## Datasheet for ABIN6699036

# Rabbit anti-Mouse IgG Antibody (DyLight 800)

# Publication



U	V	er	V	IE	!W

Durana	Marrian In C. (1181) Analih adu Duli in bat M. 000 Caniu anatad
Product Details	
Application:	Western Blotting (WB), FLISA, Fluorescence Microscopy (FM)
Conjugate:	DyLight 800
Clonality:	Polyclonal
Host:	Rabbit
Reactivity:	Mouse
Target:	IgG
Quantity:	100 μg
Overview	

Abstract:

Troduct Botano		
Purpose:	Mouse IgG (H&L) Antibody DyLight™ 800 Conjugated	
Immunogen:	Mouse IgG, whole molecule	
Isotype:	IgG	
Characteristics:	rabbit anti-Mouse IgG Antibody DyLight™ 800 conjugation, rabbit anti-Mouse IgG DyLight™800 conjugated Antibody,Anti-Mouse IgG DyLight 800 Antibody generated in rabbit detects reactivity to Mouse IgG.	
Labeling Ratio:	2.5	
Target Details		
Target:	IgG	

IgG Products

### Target Details

Target Details		
Target Type:	Antibody	
Background:	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75 % of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.	
Application Details		
Application Notes:	FLISA_Dilution: >1:20,000  IF_Microscopy_Dilution: >1:5,000  Western_Blot_Dilution: >1:10,000  Other: User Optimized	
Comment:	Anti-Mouse IgG DyLight 800 Antibody has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Conjugated Anti-Mouse IgG is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. DyLight conjugated antibodies are effective alternatives to Alexa 800 and IR Dye 800 conjugates. The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation.	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Reconstitution Volume: 100 µL  Reconstitution Buffer: Restore with deionized water (or equivalent)	
Concentration:	1.0 mg/mL	

## Handling

Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum
	Albumin (BSA) - Immunoglobulin and Protease free, 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store conjugated secondary antibody at 4° C prior to restoration. For extended storage aliquot
	contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge
	product if not completely clear after standing at room temperature. Conjugated Secondary
	Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate
	use.
Expiry Date:	12 months
Publications	
Product cited in:	Zhang, Kang, Zhou, Cui, Jia, Hu, Ji, Yuan, Cui, Shi: "Amelioratory Effects of Testosterone
	Propionate on Age-related Renal Fibrosis via Suppression of TGF-β1/Smad Signaling and
	Activation of Nrf2-ARE Signaling." in: Scientific reports, Vol. 8, Issue 1, pp. 10726, (2018) (
	PubMed).
	. 4554).