

Datasheet for ABIN6699610
CCL2 Protein



[Go to Product page](#)

2 Images

Overview

Quantity:	10 µg
Target:	CCL2
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	Mouse Monocyte Chemotactic Protein-1 (CCL2) Recombinant Protein
Purification:	Monocyte Chemotactic protein-1 (CCL2) purity was determined to be greater than 98% as determined by HpLC, analysis by UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-pAGE.
Purity:	98,00%
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/µg protein.
Biological Activity Comment:	The activity is determined by its ability to chemoattract human PBMCs or THP-1 cells at 4 - 15 ng/mL.

Target Details

Target:	CCL2
Alternative Name:	Ccl2 (CCL2 Products)
Background:	Synonyms: JE, MCAF

Target Details

Background: Monocyte Chemotactic Protein 1 (MCP-1), also known as CCL2, is thought to be produced by injured or infected tissues. MCP-1 signals through G protein-coupled receptors, CCR2 and CCR4, to recruit memory T cells, monocytes and dendritic cells. Recombinant mouse MCP-1 is a non-glycosylated protein, containing 125 amino acids, with a molecular weight of 13.8 kDa.

UniProt: [Q5SVU3](#)

Pathways: [Cellular Response to Molecule of Bacterial Origin, Positive Regulation of Immune Effector Process, ER-Nucleus Signaling, Unfolded Protein Response, The Global Phosphorylation Landscape of SARS-CoV-2 Infection](#)

Application Details

Application Notes: Other: User Optimized
Application_Note: Monocyte Chemotactic Protein-1 Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Monocyte Chemotactic Protein-1 in immunological assays.

Comment: Suggested_Applications: Cellular Assay
Other_Performance_Data:

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitution_Buffer: Restore with deionized water (or equivalent)
Reconstitution_Volume: 10 µL (10-100 µL)

Buffer: Buffer: 0.1 % Trifluoroacetic acid
Stabilizer: None

Preservative: Without preservative


Storage: 4 °C, -20 °C

Storage Comment: Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing

at room temperature.


Expiry Date: 6 months

Images



SDS-PAGE

Image 1. SDS-PAGE of Mouse Monocyte Chemotactic Protein-1 (CCL2) Recombinant Protein Bioactivity of Mouse Monocyte Chemotactic Protein-1 (CCL2) Recombinant Protein. Human THP-1 cells were allowed to migrate to Mouse MCP-1 at (0, 0.1, 1, 10, 100 and 1000 ng/mL). After 45 minutes, cells that migrated were counted using a luminescent substrate and displayed on the bar graph above. Significant increases in migration over basal levels were seen in response to Mouse MCP-1 starting at 1 ng/mL. This value is comparable to expected ranges of a chemotactic response of primary human monocytes.



SDS-PAGE

Image 2. SDS-PAGE of Mouse Monocyte Chemotactic Protein-1 (CCL2) Recombinant Protein SDS-PAGE of Mouse Monocyte Chemotactic Protein-1 (CCL2) Recombinant Protein. Lane 1: Molecular weight marker. Lane 2: 1 µg Mouse MCP-1 in non-reducing conditions . Lane 3: 1 µg Mouse MCP-1 in reducing conditions (+). Mouse MCP-1 has a predicted MW of 13.8 kDa.