

Datasheet for ABIN6699798

**EGF Protein****2** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	EGF
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

## Product Details

Purpose:	Mouse Epidermal Growth Factor Recombinant Protein
Purification:	Epidermal Growth Factor purity was determined to be greater than 97% as determined by analysis by HpLC, UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-pAGE.
Purity:	97,00%
Endotoxin Level:	Measured by LAL is typically $\leq 1$ EU/µg protein.
Biological Activity Comment:	The activity is determined by the dose-dependent proliferation of mouse BALB/c 3T3 cells and is typically less than 0.1 ng/mL.

## Target Details

Target:	EGF
Alternative Name:	Egf ( <a href="#">EGF Products</a> )
Background:	Synonyms: Urogastrone, URG Background: Epidermal Growth Factor (EGF) is a growth factor that stimulates the proliferation

## Target Details

---

of epithelial and epidermal cells. EGF family members are characterized by three intramolecular disulfide bonds and can bind to four different receptor tyrosine kinases known as EGFR/ErbB1, ErbB2, ErbB3, and ErbB4. Recombinant mouse EGF is a non-glycosylated protein, containing 54 amino acids, including 3 intra-molecular disulfide-bonds, with a molecular weight of 6.2 kDa.

UniProt: [P01132](#)

Pathways: [NF-kappaB Signaling](#), [RTK Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Regulation of Carbohydrate Metabolic Process](#), [Hepatitis C](#), [Protein targeting to Nucleus](#), [Interaction of EGFR with phospholipase C-gamma](#), [Thromboxane A2 Receptor Signaling](#), [EGFR Downregulation](#)

## Application Details

---

Application Notes: Other: User Optimized

Application\_Note: Epidermal Growth Factor Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Epidermal Growth Factor in immunological assays.

Comment: Suggested\_Applications: Cellular Assay  
Other\_Performance\_Data:

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Reconstitution\_Buffer: Restore with deionized water (or equivalent)  
Reconstitution\_Volume: 100 µL

Buffer: Buffer: 0.01 M Sodium Phosphate, pH 7.5  
Stabilizer: None

Preservative: Without preservative


Storage: 4 °C, -20 °C

Storage Comment: Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing

at room temperature.


Expiry Date: 6 months

Images



SDS-PAGE

**Image 1.** SDS-PAGE of Mouse Epidermal Growth Factor Recombinant Protein. Lane 1: 1 µg Mouse EGF in non-reducing conditions. Lane 2: 1 µg Mouse EGF in reducing conditions (+). Lane 3: Molecular weight marker. Mouse EGF is predicted to be a homodimer with a MW of 12.4 kDa.



SDS-PAGE

**Image 2.** SDS-PAGE of Mouse Epidermal Growth Factor Recombinant Protein Bioactivity of Mouse Epidermal Growth Factor Recombinant Protein. 3T3 cells were cultured with 0 to 1 ng/mL Mouse EGF. Cell proliferation was measured after 44 hours and the linear portion of the curve was used to calculate the ED50. The ED50 of Mouse EGF is 30-40 pg/mL. This value is comparable to the typical expected range less than 100 pg/mL.