

## Datasheet for ABIN6699859 **FGF4 Protein**



[Go to Product page](#)

### Overview

Quantity:	25 µg
Target:	FGF4
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

### Product Details

Purpose:	Human Fibroblast Growth Factor-4 Recombinant Protein
Purification:	Fibroblast Growth Factor-4 purity was determined to be greater than 98% as determined by analysis by reducing and non-reducing SDS-PAGE.
Purity:	98,00%
Endotoxin Level:	Measured by LAL is typically $\leq 1$ EU/µg protein.
Biological Activity Comment:	The activity is determined by its ability to induce the proliferation of mouse NR6R-3T3 fibroblasts and is typically 0.25-1.25 ng/mL.

### Target Details

Target:	FGF4
Alternative Name:	FGF4 ( <a href="#">FGF4 Products</a> )
Background:	Synonyms: Transforming protein KS3, Heparin-binding growth factor (HBGF-4), Heparin secretory-transforming protein 1 (HST-1)

## Target Details

Background: Fibroblast Growth Factor 4 (FGF-4) is a growth factor predominantly expressed during embryonic development, playing a key role in limb development. In culture, FGF-4 has been shown to be an important regulator of growth for stem cells, fibroblasts and endothelial cells. Unglycosylated FGF-4 is N-terminally cleaved into 13 kDa or 15 kDa proteins that are more active than the precursor 19 kDa protein, Bellosa P, et al. (1993). Human FGF-4 shares high homology and cross-reactivity with the mouse protein. Recombinant human FGF-4 is a non-glycosylated protein containing 140 amino acids, with a total molecular weight of 15 kDa.

UniProt: [P08620](#)

Pathways: [RTK Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Stem Cell Maintenance](#)

## Application Details

Application Notes: Other: User Optimized  
Application\_Note: Fibroblast Growth Factor-4 Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Fibroblast Growth Factor-4 in immunological assays.

Comment: Suggested\_Applications: Cellular Assay  
Other\_Performance\_Data:

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Reconstitution: Reconstitution\_Buffer: Restore with deionized water (or equivalent)  
Reconstitution\_Volume: 25 µL (25-250 µL)

Buffer: Lyophilized in 10 mM sodium phosphate, 75 mM sodium chloride, pH 7.5.

Preservative: Without preservative

Storage: 4 °C, -20 °C

Storage Comment: Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing

Handling

---

at room temperature.

---

Expiry Date: 6 months