

# Datasheet for ABIN6699860

## **FGF4 Protein**



#### Overview

Quantity:	100 μg	
Target:	FGF4	
Origin:	Human	
Source:	Escherichia coli (E. coli)	
Protein Type:	Recombinant	
Application:	SDS-PAGE (SDS)	

### **Product Details**

Purpose:	Human Fibroblast Growth Factor-4 Recombinant Protein	
Purification:	Fibroblast Growth Factor-4 purity was determined to be greater than 98% as determined by analysis by reducing and non-reducing SDS-pAGE.	
Purity:	98,00%	
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/μg protein.	
Biological Activity Comment:	The activity is determined by its ability to induce the proliferation of mouse NR6R-3T3 fibroblasts and is typically 0.25-1.25 ng/mL.	

# Target Details

Target:	FGF4	
Alternative Name:	FGF4 (FGF4 Products)	
Background:	Synonyms: Transforming protein KS3, Heparin-binding growth factor (HBGF-4), Heparin secretory-transforming protein 1 (HST-1)	

Background: Fibroblast Growth Factor 4 (FGF-4) is a growth factor predominantly expressed during embryonic development, playing a key role in limb development. In culture, FGF-4 has been shown to be an important regulator of growth for stem cells, fibroblasts and endothelial cells. Unglycosylated FGF-4 is N-terminally cleaved into 13 kDa or 15 kDa proteins that are more active than the precursor 19 kDa protein, Bellosa P, et al. (1993). Human FGF-4 shares high homology and cross-reactivity with the mouse protein. Recombinant human FGF-4 is a non-glycosylated protein containing 140 amino acids, with a total molecular weight of 15 kDa.

UniProt:

P08620

Pathways:

RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Stem Cell Maintenance

### **Application Details**

Application Notes:

Application Note: Fibroblast Growth Factor-4 Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Fibroblast Growth Factor-4 in immunological assays. Lyophilized in 10 mM sodium phosphate, 75 mM sodium chloride, pH 7.5.

Other: User Optimized

Restrictions:

For Research Use only

#### Handling

Format:	Lyophilized	
Reconstitution:	Reconstitution_Buffer: Restore with deionized water (or equivalent)  Reconstitution_Volume: 100 µL	
Buffer:	Lyophilized in 10 mM sodium phosphate, 75 mM sodium chloride, pH 7.5.	
Preservative:	Without preservative	
Storage:	4 °C,-20 °C	
Storage Comment:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.	

		- 11	١٠	
Н	ar	nd	IIn	IC

Expiry Date:

6 months