

Datasheet for ABIN6699879
Follistatin Protein (FST)



[Go to Product page](#)

2 Images

Overview

Quantity:	20 µg
Target:	Follistatin (FST)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	Human Follistatin Recombinant Protein
Purification:	Follistatin purity was determined to be greater than 95% as determined by analysis by reducing and non-reducing SDS-pAGE.
Purity:	95,00%
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/µg protein.
Biological Activity Comment:	The activity is determined by the dose-dependent neutralization of 7.5 ng/mL human Activin A. Complete neutralization is typically reached at less than 0.3 ug/mL.

Target Details

Target:	Follistatin (FST)
Alternative Name:	FST (FST Products)
Background:	Synonyms: FS, activin-binding protein, FSH-suppressing protein (FSP) Background: Follistatin is an autocrine acting protein that is expressed by many tissues, but at

Target Details

notably higher levels in the ovary and skin. Follistatin functions to negatively regulate the signaling of a wide variety of TGF-beta family members (activin, BMPs, myostatin, GDF-11 and TGF-beta 1). Mechanistically, follistatin works as an antagonist by complexing with TGF-beta family members to prevent them from interacting with their signaling receptors. Recombinant human Follistatin is a non-glycosylated protein, containing 289 amino acids, with a molecular weight of 31.7 kDa.

UniProt: [P19883](#)

Pathways: [Negative Regulation of Hormone Secretion](#)

Application Details

Application Notes: Other: User Optimized
Application_Note: Follistatin Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Follistatin in immunological assays.

Comment: Suggested_Applications: Cellular Assay
Other_Performance_Data:

Restrictions: For Research Use only

Handling

Format: Lyophilized


Reconstitution: Reconstitution_Buffer: Restore with deionized water (or equivalent)
Reconstitution_Volume: 20 µL (20-200 µL)

Buffer: This product is provided as a lyophilized powder, in 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5.

Preservative: Without preservative


Storage: 4 °C, -20 °C

Storage Comment: Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.



SDS-PAGE

Image 1. SDS-PAGE of Human Follistatin Recombinant Protein. Lane 1: Molecular weight marker. Lane 2: 1 µg Human Follistatin in non-reducing conditions. Lane 3: 1 µg Human Follistatin in reducing conditions (+). Human Follistatin has a predicted MW of 31.7 kDa.



SDS-PAGE

Image 2. SDS-PAGE of Human Follistatin Recombinant Protein. Bioactivity of Human Follistatin Recombinant Protein. MPC-11 cells were cultured with 0 to 1 µg/mL Human Follistatin with or without 7.5 ng/mL Human Activin A. Cell viability was measured after 66 hours and the linear portion of the curve was used to calculate the ED50. The ED50 of Human Follistatin is 30-50 ng/mL. Complete neutralization is seen by 250 ng/mL.