

Datasheet for ABIN6699880  
**Follistatin Protein (FST)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	100 µg
Target:	Follistatin (FST)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

## Product Details

Purpose:	Human Follistatin Recombinant Protein
Purification:	Follistatin purity was determined to be greater than 95% as determined by analysis by reducing and non-reducing SDS-pAGE.
Purity:	95,00%
Endotoxin Level:	Measured by LAL is typically $\leq 1$ EU/µg protein.
Biological Activity Comment:	The activity is determined by the dose-dependent neutralization of 7.5 ng/mL human Activin A. Complete neutralization is typically reached at less than 0.3 ug/mL.

## Target Details

Target:	Follistatin (FST)
Alternative Name:	FST ( <a href="#">FST Products</a> )
Background:	Synonyms: FS, activin-binding protein, FSH-suppressing protein (FSP) Background: Follistatin is an autocrine acting protein that is expressed by many tissues, but at

## Target Details

---

notably higher levels in the ovary and skin. Follistatin functions to negatively regulate the signaling of a wide variety of TGF-beta family members (activin, BMPs, myostatin, GDF-11 and TGF-beta 1). Mechanistically, follistatin works as an antagonist by complexing with TGF-beta family members to prevent them from interacting with their signaling receptors. Recombinant human Follistatin is a non-glycosylated protein, containing 289 amino acids, with a molecular weight of 31.7 kDa.

UniProt: [P19883](#)

Pathways: [Negative Regulation of Hormone Secretion](#)

## Application Details

---

Application Notes: Other: User Optimized  
Application\_Note: Follistatin Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Follistatin in immunological assays.

Comment: Suggested\_Applications: Cellular Assay  
Other\_Performance\_Data:

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized


Reconstitution: Reconstitution\_Buffer: Restore with deionized water (or equivalent)  
Reconstitution\_Volume: 100 µL

Buffer: This product is provided as a lyophilized powder, in 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5.

Preservative: Without preservative


Storage: 4 °C, -20 °C

Storage Comment: Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.



### SDS-PAGE

**Image 1.** SDS-PAGE of Human Follistatin Recombinant Protein. Lane 1: Molecular weight marker. Lane 2: 1 µg Human Follistatin in non-reducing conditions. Lane 3: 1 µg Human Follistatin in reducing conditions (+). Human Follistatin has a predicted MW of 31.7 kDa.



### SDS-PAGE

**Image 2.** SDS-PAGE of Human Follistatin Recombinant Protein. Bioactivity of Human Follistatin Recombinant Protein. MPC-11 cells were cultured with 0 to 1 µg/mL Human Follistatin with or without 7.5 ng/mL Human Activin A. Cell viability was measured after 66 hours and the linear portion of the curve was used to calculate the ED50. The ED50 of Human Follistatin is 30-50 ng/mL. Complete neutralization is seen by 250 ng/mL.