

Datasheet for ABIN6699899

GM-CSF Protein[Go to Product page](#)

Overview

Quantity:	20 µg
Target:	GM-CSF (CSF2)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	Human Granulocyte Macrophage Colony Stimulating Factor Recombinant Protein (Animal Free)
Purification:	Granulocyte Macrophage Colony Stimulating Factor is produced with no animal-derived raw products, animal free equipment and animal free protocols. Purity was determined to be greater than 95% as determined by analysis by UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-PAGE.
Purity:	95,00%
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/µg protein.
Grade:	Animal-Free
Biological Activity Comment:	The activity is determined by the dose-dependent induction of human TF-1 cell proliferation and is typically less than 0.1 ng/mL.

Target Details

Target:	GM-CSF (CSF2)
---------	---------------

Target Details

Alternative Name:	CSF2 (CSF2 Products)
Background:	Synonyms: Molgramostin, Pluripoietin-α, MGI1GM, Sargramostim Background: Granulocyte Macrophage Colony Stimulating Factor (GM-CSF) is hematopoietic factor produced by endothelial cells, monocytes, fibroblasts and T cells in response to a number of inflammatory mediators. GM-CSF is able to stimulate the production of neutrophilic granulocytes, macrophages, and mixed granulocyte-macrophage colonies from bone marrow cells. GM-CSF can also stimulate some functional activities in mature granulocytes and macrophages. Human and mouse GM-CSF show no cross-reactivity. Recombinant human GM-CSF is a non-glycosylated protein, containing 128 amino acids, with a molecular weight of 14.6 kDa.
UniProt:	P04141
Pathways:	JAK-STAT Signaling , Cellular Response to Molecule of Bacterial Origin

Application Details

Application Notes:	Application Note: Granulocyte Macrophage Colony Stimulating Factor Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Granulocyte Macrophage Colony Stimulating Factor in immunological assays. Other: User Optimized
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Reconstitution_Buffer: Restore with deionized water (or equivalent) Reconstitution_Volume: 20 μL (20-200 μL)
Buffer:	Buffer: 0.01 M Sodium Phosphate, pH 7.5 Stabilizer: None
Preservative:	Without preservative
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each

Handling

opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.

Expiry Date: 6 months