antibodies -online.com





HSD17B10 Protein (GST tag)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	50 μg
Target:	HSD17B10
Origin:	Human
Source:	Baculovirus infected Insect Cells, Insect cells (Sf9)
Purification tag / Conjugate:	This HSD17B10 protein is labelled with GST tag.
Application:	Western Blotting (WB)

Product Details

Cross-Reactivity:	Human
Characteristics:	Hydroxysteroid (17-beta) Dehydrogenase 10, ABAD, ERAB, HCD2, MHBD, HADH2, SCHAD, 17b-
	HSD10, CAMR, MRPP2, MRX17, MRX31, MRXS10, SDR5C1, DUPXp11.22, 3-hydroxyacyl-CoA
	dehydrogenase type-2, 17-beta-hydroxysteroid dehydrogenase 10, 3-hydroxy-2-methylbutyryl-
	CoA dehydrogenase, 3-hydroxyacyl-CoA dehydrogenase type II, Endoplasmic reticulum-
	associated amyloid beta-peptide-binding protein, Mitochondrial ribonuclease P protein 2, Short-
	chain type dehydrogenase/reductase XH98G2, Type II HADH
Purification:	Recombinant full-length human HSD17B10 was expressed using baculovirus in Sf9 cells using
	an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >95%
	by densitometry.

Target Details

Target:	HSD17B10	
Alternative Name:	HSD17B10 (HSD17B10 Products)	

Application Details

Expiry Date:

12 months

Application Notes:	Application Note: HSD17B10 Protein is stored in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM		
	glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, 25 % glycerol. HSD17B10 Protein is		
	suitable for use in Western Blot. Expect a band approximately $\sim\!51~\text{kDa}$ on specific lysates or		
	tissues. Specific conditions for reactivity should be optimized by the end user.		
	Western Blot Dilution: User Optimized		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	0.2 μg/μL		
Storage:	-80 °C		
Storage Comment:	-80 °c		