

Datasheet for ABIN6700048

IL16 Protein**2** Images[Go to Product page](#)

Overview

Quantity:	10 µg
Target:	IL16
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	Mouse Interleukin-16 Recombinant Protein
Purification:	Interleukin-16 purity was determined to be greater than 95% as determined by HPLC, analysis by UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-pAGE.
Purity:	95,00%
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/µg protein.
Biological Activity Comment:	The activity is determined by the ability to chemoattract primary human T cells and is typically less than 200 ng/mL.

Target Details

Target:	IL16
Alternative Name:	Il16 (IL16 Products)
Background:	Synonyms: Lymphocyte Chemoattractant Factor (LCF) Background: Interleukin 16 (IL-16) is produced primarily by CD4+ and CD8+ T cells and acts as

Target Details

a chemo-attractant for lymphocytes, monocytes, eosinophils, dendritic cells and Langerhans cells. Additionally, IL-16 has been reported to upregulate IL-2 receptor (CD25), induce progression of cells to the G1 phase and suppress HIV & SIV replication. Recombinant mouse IL-16 is a non-glycosylated protein, containing 127 amino acids, with a molecular weight of 13.2 kDa.

UniProt: [O54824](#)

Application Details

Application Notes: Other: User Optimized
Application_Note: Interleukin-16 Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Interleukin-16 in immunological assays.

Comment: Suggested_Applications: Cellular Assay
Other_Performance_Data:

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitution_Buffer: Restore with deionized water (or equivalent)
Reconstitution_Volume: 10 μ L (10-100 μ L)



Concentration: 0.1 mg/mL

Buffer: Buffer: 0.01 M Sodium Phosphate, pH 7.5
Stabilizer: None

Preservative: Without preservative

Storage: 4 °C, -20 °C

Storage Comment: Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.



SDS-PAGE

Image 1. SDS-PAGE of Mouse Interleukin-16 Recombinant Protein Bioactivity of Mouse Interleukin-16 Recombinant Protein. Human T cells were allowed to migrate to Mouse IL-16 at (0, 5, 20, 100 and 500 ng/mL). After 4 hours, cells that migrated were counted using a luminescent substrate and displayed on the bar graph above. Significant increases in migration over basal levels were seen in response to Mouse IL-16 starting at 5 ng/mL.

SDS-PAGE

Image 2. SDS-PAGE of Mouse Interleukin-16 Recombinant Protein SDS-PAGE of Mouse Interleukin-16 Recombinant Protein. Lane 1: 1 µg Mouse IL-16 in non-reducing conditions. Lane 2: 1 µg Mouse IL-16 in reducing conditions (+). Lane 3: Molecular weight marker. Mouse IL-16 has a predicted MW of 13.2 kDa.