

Datasheet for ABIN6700065

IL-2 Protein



Overview

Quantity:	50 μg
Target:	IL-2 (IL2)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	Human Interleukin-2 Recombinant Protein (Animal Free)
Purification:	Interleukin-2 is produced with no animal-derived raw products, animal free equipment and animal free protocols. Purity was determined to be greater than 90% as determined by analysis by UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-PAGE.
Purity:	90,00%
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/μg protein.
Grade:	Animal-Free
Biological Activity Comment:	The activity is measured by the dose-dependent proliferation of CTLL-2 cells and is typically less than 1 ng/mL.

Target Details

Target:	IL-2 (IL2)
Alternative Name:	IL2 (IL2 Products)

Target Details

Background:	Synonyms: T cell growth factor, TCGF, Aldesleukin
	Background: Interleukin-2 (IL-2) is an immuno-modulatory cytokine produced by stimulated T
	cells. IL-2 signals through the IL-2R receptor to induce proliferation of activated T cells. IL-2 also
	stimulates proliferation and differentiation of B cells, natural killer cells, monocytes and
	macrophages. Recombinant human IL-2 is a non-glycosylated protein, containing 134 amino
	acids, with a molecular weight of 15.5 kDa. Recombinant human IL-2 has a Ser substitute for
	Cysteine at position 126.
UniProt:	P60568
Pathways:	JAK-STAT Signaling, Regulation of Leukocyte Mediated Immunity, Positive Regulation of
	Immune Effector Process, Production of Molecular Mediator of Immune Response, Activated T
	Cell Proliferation
Application Details	
Application Notes:	Application Note: Human Interleukin-2 Recombinant Protein is suitable as a control for
- In the control of t	polyclonal or monoclonal anti-Human Interleukin-2 in immunological assays.
	Other: User Optimized
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Reconstitution_Buffer: Restore with deionized water (or equivalent)
	Reconstitution_Volume: 50µL
Concentration:	0.1 mg/mL
Buffer:	Buffer: 0.1 % Trifluoroacetic acid
	Stabilizer: None
Preservative:	Without preservative
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This
	product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier
	protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and
	freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each
	opening to dislodge contents from the cap and to clarify if contents are not clear after standing

Handling	
----------	--

at room temperature.

Expiry Date: 6 months