

Datasheet for ABIN6700385

PDGFA Protein



_			
()	V/C	rv	٨/

Quantity:	100 μg
Target:	PDGFA
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	Human Platelet Derived Growth Factor-AA Recombinant Protein (Animal Free)	
Purification:	Platelet-Derived Growth Factor is produced with no animal-derived raw products, animal free equipment and animal free protocols. Purity was determined to be greater than 97% as determined by analysis by UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-PAGE.	
Purity:	97,00%	
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/µg protein.	
Grade:	Animal-Free	
Biological Activity Comment:	The activity is determined by the dose-dependent proliferation of mouse 3T3 indicator cells and is typically 3-5 ng/mL.	

Target Details

Target:	PDGFA

Target Details

Alternative Name:	PDGFA (PDGFA Products)	
Background:	Synonyms: PDGF-1, Platelet-derived growth factor A chain, Platelet-derived growth factor alpha	
	polypeptide	
	Background: Platelet-Derived Growth Factor (PDGF) is a mitogenic peptide growth hormone	
	carried in the alpha-granules of platelets and is released when platelets adhere to traumatized	
	tissues. Connective tissue cells near the traumatized region respond by initiating the process o	
	replication. The synthesis of PDGF can be induced by IL-1, IL-6, TNF- α , TGF- β and EGF.	
	Recombinant human PDGF-AA is a non-glycosylated disulfide-linked homodimer, containing	
	two 125 amino acid chains, with a total molecular weight of 28.5 kDa.	
UniProt:	P04085	
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin	
	Signaling Pathway, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor	
	Signaling	
Application Details		
	Application Nata Distalat Darived Crowth Factor AA Decembinant Protein is quitable as a	
Application Notes:	Application Note: Platelet Derived Growth Factor-AA Recombinant Protein is suitable as a	
	control for polyclonal or monoclonal anti-Platelet Derived Growth Factor-AA in immunological assays.	
	Other: User Optimized	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Reconstitution_Buffer: Restore with deionized water (or equivalent)	
	Reconstitution_Volume: 100 μL	
Concentration:	0.1 mg/mL	
Buffer:	Buffer: 0.1 % Trifluoroacetic acid	
	Stabilizer: None	
Preservative:	Without preservative	
Storage:	4 °C,-20 °C	
Storage Comment:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This	

Handling

product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.

Expiry Date:

6 months