

# Datasheet for ABIN6700565

## **STAT1 Protein (GST tag)**



| _ |       |             |    |    |             |     |
|---|-------|-------------|----|----|-------------|-----|
|   | V     | $\triangle$ | r۱ | /1 | $\triangle$ | Λ/  |
|   | ' V ' |             | ΙV |    |             | v v |

| Quantity:                     | 20 μg  |
|-------------------------------|--|
| Target:                       | STAT1  |
| Origin:                       | Human  |
| Source:                       | Insect cells (Sf9)                           |
| Protein Type:                 | Recombinant                                  |
| Purification tag / Conjugate: | This STAT1 protein is labelled with GST tag. |
| Application:                  | Western Blotting (WB)                        |

### **Product Details**

| Purpose:      | STAT1 beta recombinant protein-GST fusion protein  |
|---------------|--|
| Purification: | Recombinant full-length human STAT1 $\beta$ was expressed by baculovirus in Sf9 insect cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >90% by densitometry. |
| Purity:       | >90%   |

#### **Target Details**

| Target:           | STAT1   |  |
|-------------------|---|--|
| Alternative Name: | STAT1 (STAT1 Products)  |  |
| Background:       | Synonyms: ISGF-3, STAT91, DKFZp686B04100, Signal transducer and activator of transcription 1-alpha/beta, Transcription factor ISGF-3 components p91/p84 |  |
|                   | Background: STAT1β is a member of the signal transducers and activators of transcription  |  |

(STAT) family of proteins that carry out a dual function: signal transduction and activation of transcription. STAT1 $\beta$  transcription factor is specific for the IFN pathway and plays a central role in mediating many, if not all, IFN-dependent biological responses (1). Presence of STAT1 $\beta$  leads to an efficient antiviral response when cells were infected with virus suggesting that a STAT-dependent pathway is activated following virus infection by endogenously produced IFN. Virus-induced STAT protein translocation from the cytoplasmic compartment can be detected within 3 h of infection. STAT1 $\beta$  Protein is ideal for investigators involved in Signaling Proteins, Transcription Proteins, Apoptosis/Autophagy, Cancer, ERK/MAPK Pathway, Inflammation, and JAK/STAT Pathway research.

NCBI Accession:

NM\_139266

Pathways:

JAK-STAT Signaling, RTK Signaling, Interferon-gamma Pathway, Response to Growth Hormone Stimulus, Cellular Response to Molecule of Bacterial Origin, Positive Regulation of Endopeptidase Activity, Hepatitis C, CXCR4-mediated Signaling Events

#### **Application Details**

Application Notes:

Western\_Blot\_Dilution: User Optimized

Other: Kinase Assay-User Optimized

Application\_Note: STAT1 beta Protein is suitable for use in Western Blot and Kinase Assay.

Expect a band approximately ~118 kDa on specific lysates or tissues. Specific conditions for

reactivity should be optimized by the end user.

Restrictions:

For Research Use only

#### Handling

| Format:          | Liquid  |
|------------------|---|
| Concentration:   | 0.2 μg/μL   |
| Buffer:          | STAT1 beta Protein is stored in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, 25 % glycerol.   |
| Storage:         | -80 °C  |
| Storage Comment: | Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles. |
| Expiry Date:     | 12 months   |