

Datasheet for ABIN6700669

**MAPT Protein (Ser214Ala-Mutant)**[Go to Product page](#)

## Overview

Quantity:	20 µg
Target:	MAPT
Protein Characteristics:	Ser214Ala-Mutant
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	Western Blotting (WB)

## Product Details

Purpose:	Tau-441 (S214A) recombinant protein
Purification:	Recombinant human tag-free Tau-441 (S214A) was expressed in E. coli cells. The purity was determined to be >90% by densitometry.
Purity:	>90%

## Target Details

Target:	MAPT
Alternative Name:	MAPT ( <a href="#">MAPT Products</a> )
Background:	Synonyms: Tau-F, (N2R4), Tau-4, MAPT, MSTD, PPND, DDPAC, MAPTL, MTBT1, MTBT2, FTDP-17, FLJ31424, MGC138549, Microtubule-associated protein tau Background: Tau-441 or Tau-F is a member of the Tau family of proteins which function to stabilize the microtubules by binding to them. Tau proteins are subject to phosphorylation and

## Target Details

---

this phenomenon regulates the association of the Tau protein with the microtubules (1). Deposits of Alzheimer's disease AD-associated proteins, such as hyperphosphorylated Tau, as well as other shared misfolded proteins, such as,  $\beta$ -amyloid precursor protein ( $\beta$ APP), ubiquitin, and various chaperones and protein kinases are thought to play a pathologic role in the cognitive decline and muscular failure. Malfunctioning of Tau proteins is associated with microtubules disintegration and collapsing of the neuronal transport system (2). Tau-441 Protein is ideal for investigators involved in Signaling Proteins, Tau Proteins, Invasion/Metastasis, Neurobiology, and p38 Pathway research.

UniProt: [P10636-8](#)

Pathways: [MAPK Signaling](#), [Microtubule Dynamics](#), [M Phase](#), [Regulation of Cell Size](#)

## Application Details

---

Application Notes: Western\_Blot\_Dilution: User Optimized  
Other: Kinase Assay-User Optimized  
Application\_Note: Tau-441 Protein is suitable for use in Western Blot and Kinase Assay. Expect a band approximately ~64 kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 0.2  $\mu$ g/ $\mu$ L

Buffer: Tau-441 Protein is stored in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.25 mM DTT, 0.1 mM PMSF, 25 % glycerol.

Storage: -80  $^{\circ}$ C

Storage Comment: Store product at -70 $^{\circ}$ C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

Expiry Date: 12 months