

### Datasheet for ABIN6700784

# 14-3-3 zeta Protein (YWHAZ)



#### Overview

OVEIVIEW	
Quantity:	20 μg
Target:	14-3-3 zeta (YWHAZ)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	Western Blotting (WB)
Product Details	
_	

Purpose:	14-3-3 zeta recombinant protein
Purification:	Recombinant full-length human 14-3-3 $\zeta$ was expressed in E. coli cells. The purity was determined to be >95% by densitometry.
Purity:	>95%

## Target Details

Target:	14-3-3 zeta (YWHAZ)
Alternative Name:	YWHAZ (YWHAZ Products)
Background:	Synonyms: 14-3-3 zeta, YWHAZ, KCIP-1, MGC111427, MGC126532, MGC138156, 14-3-3 protein zeta, 14-3-3-like protein, Protein Leonardo, 14-3-3, 14-3-3EZ, leo, THAP
	Background: 14-3-3ζ (also known as tyrosine 3-monooxygenase/ tryptophan 5-monooxygenase
	activation protein, zeta polypeptide) is a member of the 14-3-3 family of proteins which mediate
	signal transduction by binding to phosphoserine-containing proteins. 14-3-3 $\zeta$ protein plays a

key role in cancer biology by being an important regulator of major cellular processes such as proliferation, differentiation, senescence and apoptosis (1). 14-3-3ζ protein has been shown to interact with the IRS1 protein, suggesting a role for this protein in regulating insulin sensitivity by interrupting the association between the insulin receptor and IRS1 (2). 14-3-3ζ Protein is ideal for investigators involved in Cell Stress& Chaperone Proteins, Cell Signaling, AKT/PKB Pathway, Cancer, Cell Cycle, Cellular Stress, ERK/MAPK Pathway, Neurobiology, PKA/PKC Pathway, WNT Signaling research.

NCBI Accession:

NM\_003406

Pathways:

Apoptosis, Hormone Transport, Myometrial Relaxation and Contraction, Regulation of
Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Synaptic
Membrane, Production of Molecular Mediator of Immune Response, Maintenance of Protein
Location

### **Application Details**

Application Notes:

Western\_Blot\_Dilution: User Optimized

Application\_Note: 14-3-3 $\zeta$  Protein is suitable for use in Western Blot. Expect a band approximately ~ 29 kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.

Restrictions:

For Research Use only

### Handling

Format:	Liquid
Concentration:	0.2 μg/μL
Buffer:	14-3-3ζ Protein is stored in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, 25 % glycerol.
Storage:	-80 °C
Storage Comment:	Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
Expiry Date:	12 months