

Datasheet for ABIN6700855 **VEGFA Protein**



[Go to Product page](#)

Overview

Quantity:	10 µg
Target:	VEGFA
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

Product Details

Purpose:	Human Vascular Endothelial Growth Factor-121 Recombinant Protein (Animal Free)
Purification:	Vascular Endothelial Growth Factor is produced with no animal-derived raw products, animal free equipment and animal free protocols. Purity was determined to be greater than 98% as determined by reducing and non-reducing SDS-PAGE.
Purity:	98,00%
Endotoxin Level:	Measured by LAL is typically ≤ 1 EU/µg protein.
Grade:	Animal-Free
Biological Activity Comment:	The activity is determined by the dose-dependent proliferation of HUVECs and is typically 5 ng/mL.

Target Details

Target:	VEGFA
Alternative Name:	VEGFA (VEGFA Products)

Target Details

Background:	Synonyms: VEGF-A, glioma-derived endothelial cell mitogen, Vascular permeability factor (VPF) Background: Vascular Endothelial Growth Factor-A (VEGF-A) was originally isolated from tumor cells and is produced by a wide variety of cell types. In addition to stimulating vascular growth and vascular permeability, VEGF-A may play a role in stimulating vasodilation via nitric oxide-dependent pathways. VEGF-A has several variants, one being VEGF-121. Rat and bovine VEGF are one amino acid shorter than the human factor, and the bovine and human sequences show a homology of 95 % . Recombinant human VEGF-121 is a non-glycosylated homodimer, containing two 121 amino acid chains , with a total molecular weight of 28.4 kDa.
UniProt:	P15692-9
Pathways:	RTK Signaling , Glycosaminoglycan Metabolic Process , Regulation of Cell Size , Tube Formation , Signaling Events mediated by VEGFR1 and VEGFR2 , Platelet-derived growth Factor Receptor Signaling , VEGFR1 Specific Signals , VEGF Signaling

Application Details

Application Notes:	Application Note: Endothelial Growth Factor-121 Recombinant Protein has been tested by SDS-PAGE and is suitable as a control for polyclonal or monoclonal anti-Endothelial Growth Factor-121 in immunological assays. Other: User Optimized
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Reconstitution_Buffer: Restore with deionized water (or equivalent) Reconstitution_Volume: 10 µL (10-100 µL)
Buffer:	Buffer: 0.1 % Trifluoroacetic acid Stabilizer: None
Preservative:	Without preservative
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each

Handling

opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.

Expiry Date: 6 months