

Datasheet for ABIN6700926

**GDNF Protein****2** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	GDNF
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	SDS-PAGE (SDS)

## Product Details

Purpose:	Human Glial Derived Neurotrophic Factor Recombinant Protein
Purification:	Glial Derived Neurotrophic Factor purity was determined to be greater than 98% as determined by analysis by HpLC, UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-pAGE.
Purity:	98,00%
Endotoxin Level:	Measured by LAL is typically $\leq 1$ EU/µg protein.
Biological Activity Comment:	The activity is determined by the dose-dependent proliferation of C6 cells and is typically 0.5-5 µg/mL.

## Target Details

Target:	GDNF
Alternative Name:	GDNF ( <a href="#">GDNF Products</a> )
Background:	Synonyms: ATF-1 Background: Glial Cell Line-Derived Neurotrophic Factor (GDNF) is a neurotrophic factor that is

## Target Details

---

closely related to other neurotrophic factors, such as Neurturin, Persephin, and Artemin, by a common structural feature called the cysteine-knot. GDNF signals through a multicomponent system of receptors that includes RET and GFR $\alpha$ 1-4, to promote dopamine uptake, survival and differentiation of neurons. Recombinant human GDNF is a non-glycosylated homodimer, containing two 135 amino acid chains, with a total molecular weight of 30.4 kDa.

UniProt: [P39905](#)

Pathways: [RTK Signaling](#), [Synaptic Membrane](#), [Tube Formation](#), [Autophagy](#), [Smooth Muscle Cell Migration](#)

## Application Details

---

Application Notes: Other: User Optimized  
Application\_Note: Glial Derived Neurotrophic Factor Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Glial Derived Neurotrophic Factor in immunological assays.

Comment: Suggested\_Applications: Cellular Assay  
Other\_Performance\_Data:

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Reconstitution\_Buffer: Restore with deionized water (or equivalent)  
Reconstitution\_Volume: 100  $\mu$ L


Buffer: Lyophilized in 10 mM sodium citrate, 100 mM sodium chloride, pH 4.0.

Preservative: Without preservative

Storage: 4 °C, -20 °C


Storage Comment: Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.

Expiry Date: 6 months



### SDS-PAGE

**Image 1.** SDS-PAGE of Human Glial Derived Neurotrophic Factor Recombinant Protein Bioactivity of Human Glial Derived Neurotrophic Factor Recombinant Protein. C6 cells were cultured with 0 to 10 ug/mL Human GDNF. Cell proliferation was measured after 7 days and the linear portion of the curve was used to calculate the ED50. The ED50 of Human GDNF is 1.7-2.6 ug/mL.



### SDS-PAGE

**Image 2.** SDS-PAGE of Human Glial Derived Neurotrophic Factor Recombinant Protein SDS-PAGE of Human Human Glial Derived Neurotrophic Factor Recombinant Protein. Lane 1: 1 µg Human GDNF in non-reducing conditions. Lane 2: 1 µg Human GDNF in reducing conditions (+). Lane 3: Molecular weight marker. Human GDNF is predicted to be a disulfide linked homodimer having a total MW of 30.4 kDa.