antibodies -online.com







anti-CDK8 antibody



Publication



Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Quantity:	100 μL
Target:	CDK8
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDK8 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CDK8
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	CDK8	
Alternative Name:	CDK8 (CDK8 Products)	
Background:	Synonyms: Cyclin Dependent Kinase 8, Cell Division Kinase 8, Cell division protein kinase 8, K35	
	Mediator complex subunit CDK8, Mediator of RNA polymerase II transcription subunit CDK8, Protein kinase K35, CDK8_HUMAN, Cyclin-dependent kinase 8,.	
	Total Kildse Koo, obko_flow, w, oyelin dependent kildse o,.	

Background: The protein encoded by this gene is a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are highly similar to the gene products of Saccharomyces cerevisiae cdc28, and Schizosaccharomyces pombe cdc2, and are known to be important regulators of cell cycle progression. This kinase and its regulatory subunit cyclin C are components of the RNA polymerase II holoenzyme complex, which phosphorylates the carboxy-terminal domain (CTD) of the largest subunit of RNA polymerase II. This kinase has also been shown to regulate transcription by targeting the CDK7/cyclin H subunits of the general transcription initiation factor IIH (TFIIH), thus providing a link between the 'Mediator-like' protein complexes and the basal transcription machinery. [provided by RefSeq, Jul 2008].

Gene ID:

1024

Pathways:

Cell Division Cycle, Regulation of Lipid Metabolism by PPARalpha

Application Details

Application Notes: WB(1:100-1000), IHC-P(1:100-500), IF(IHC-P)(1:50-200)

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 1 % BSA, 50 % glycerol and 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months

Publications

Product cited in:

Li, Tian, Ren, Chen, Wang, Ge, Wu, Sun, Liu, Xiao: "MicroRNA-101 is a potential prognostic indicator of laryngeal squamous cell carcinoma and modulates CDK8." in: **Journal of translational medicine**, Vol. 13, pp. 271, (2015) (PubMed).