## antibodies -online.com





## anti-DVL1 antibody (AA 21-100) (Biotin)



Go to Product page

( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	DVL1	
Binding Specificity:	AA 21-100	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DVL1 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DVL1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	DVL1	
Alternative Name:	DVL1 (DVL1 Products)	
Background:	Synonyms: DVL, DVL1L1, DVL1P1, Segment polarity protein dishevelled homolog DVL-1,	

Target Details	
	Dishevelled-1, DSH homolog 1, DVL1
	Background: Participates in Wnt signaling by binding to the cytoplasmic C-terminus of frizzled
	family members and transducing the Wnt signal to down-stream effectors. Plays a role both in
	canonical and non-canonical Wnt signaling. Plays a role in the signal transduction pathways
	mediated by multiple Wnt genes. Required for LEF1 activation upon WNT1 and WNT3A
	signaling. DVL1 and PAK1 form a ternary complex with MUSK which is important for MUSK-
	dependent regulation of AChR clustering during the formation of the neuromuscular junction
	(NMJ).
Gene ID:	1855
UniProt:	014640
Pathways:	WNT Signaling, Synaptic Membrane, Skeletal Muscle Fiber Development
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 an

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	