

## Datasheet for ABIN671625

# anti-HMGB1 antibody (AA 75-170) (HRP)



_			
	IVe	rv	iew

Quantity:	100 μL	
Target:	HMGB1	
Binding Specificity:	AA 75-170	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HMGB1 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
5		

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human HMGB1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow
Purification:	Purified by Protein A.

### Target Details

Target:	HMGB1
Alternative Name:	HMGB1 (HMGB1 Products)

#### **Target Details**

Background:	Synonyms: HMG1, HMG3, SBP-1, High mobility group protein B1, High mobility group protein 1, HMG-1, HMGB1	
	Background: DNA binding proteins that associates with chromatin and has the ability to bend	
	DNA. Binds preferentially single-stranded DNA. Involved in V(D)J recombination by acting as a	
	cofactor of the RAG complex. Acts by stimulating cleavage and RAG protein binding at the 23	
	bp spacer of conserved recombination signal sequences (RSS). Heparin-binding protein that	
	has a role in the extension of neurite-type cytoplasmic processes in developing cells (By	
	similarity).	
Gene ID:	3146	
UniProt:	P09429	
Pathways:	p53 Signaling, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development,	
	Positive Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process,	
	Toll-Like Receptors Cascades, Smooth Muscle Cell Migration, Inflammasome	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish	
	peroxidase.	
Storage:	-20 °C	

### Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months