Datasheet for ABIN671638 anti-PIWIL1 antibody (AA 551-650) (FITC)

-online.com antibodies



~	
$() \vee \Box$	rview
Ove	

Quantity:	100 µL
Target:	PIWIL1
Binding Specificity:	AA 551-650
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIWIL1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from chicken PIWIL1
Isotype:	lgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	PIWIL1
Alternative Name:	PIWIL1 (PIWIL1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN671638 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

Background:   Synonyms: PIWI, Piwi-like protein 1, PIWIL1     Background:   Background: Plays a central role during gametogenesis by repressing transposable e     and preventing their mobilization, which is essential for the germline integrity. Acts v   piRNA metabolic process, which mediates the repression of transposable elements -     meiosis by forming complexes composed of piRNAs and Piwi proteins and govern timethylation and subsequent repression of transposons. Directly binds methylated piclass of 24 to 3 nucleotide RNAs that are generated by a Dicer-independent mechani primarily derived from transposons and other repeated sequence elements. Besides function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).     Gene ID:   416804     UniProt:   A6N7Y9     Application Details   IF(IHC-P) 1:50-200     IF(IHC-F) 1:50-200   IF(IHC-F) 1:50-200     IF(ICC) 1:50-200   IF(ICC) 1:50-200     Handling   IIquid	ia the during ne RNAs, a ism and are their
and preventing their mobilization, which is essential for the germline integrity. Acts v piRNA metabolic process, which mediates the repression of transposable elements o meiosis by forming complexes composed of piRNAs and Piwi proteins and govern the methylation and subsequent repression of transposons. Directly binds methylated pi class of 24 to 3 nucleotide RNAs that are generated by a Dicer-independent mechani primarily derived from transposons and other repeated sequence elements. Besides function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).Gene ID:416804UniProt:A6N7Y9Application DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingEventy Use only	ia the during ne RNAs, a ism and are their
piRNA metabolic process, which mediates the repression of transposable elements i meiosis by forming complexes composed of piRNAs and Piwi proteins and govern th methylation and subsequent repression of transposons. Directly binds methylated pi class of 24 to 3 nucleotide RNAs that are generated by a Dicer-independent mechani primarily derived from transposons and other repeated sequence elements. Besides function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).Gene ID:416804UniProt:A6N7Y9Application DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingHandling	during ne IRNAs, a ism and are their
meiosis by forming complexes composed of piRNAs and Piwi proteins and govern the methylation and subsequent repression of transposons. Directly binds methylated pi class of 24 to 3 nucleotide RNAs that are generated by a Dicer-independent mechani primarily derived from transposons and other repeated sequence elements. Besides function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).Gene ID:416804UniProt:A6N7Y9Application DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingHandling	ne IRNAs, a ism and are their
methylation and subsequent repression of transposons. Directly binds methylated piclass of 24 to 3 nucleotide RNAs that are generated by a Dicer-independent mechaniprimarily derived from transposons and other repeated sequence elements. Besides function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).     Gene ID:   416804     UniProt:   A6N7Y9     Application Details   IF(IHC-P) 1:50-200     IF(IHC-F) 1:50-200   IF(IHC-F) 1:50-200     IF(ICC) 1:50-200   IF(ICC) 1:50-200     IF(ICC) 1:50-200   IF(IHC-F) 1:50-200     IF(IHC-F) 1:50-200   IF(IHC-F) 1:50-200	RNAs, a ism and are their
class of 24 to 3 nucleotide RNAs that are generated by a Dicer-independent mechani primarily derived from transposons and other repeated sequence elements. Besides function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).Gene ID:416804UniProt:A6N7Y9Application DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandling	ism and are their
primarily derived from transposons and other repeated sequence elements. Besides function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).Gene ID:416804UniProt:A6N7Y9Application DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 	their
function in transposable elements repression, piRNAs are probably involved in other during meiosis such as translation regulation (By similarity).Gene ID:416804UniProt:A6N7Y9Application DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandling	
during meiosis such as translation regulation (By similarity).Gene ID:416804UniProt:A6N7Y9Application DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 FOR Research Use onlyRestrictions:For Research Use onlyHandlingImage: Comparison of the second secon	processes
Gene ID:   416804     UniProt:   A6N7Y9     Application Details   IF(IHC-P) 1:50-200     IF(IHC-F) 1:50-200   IF(IHC-F) 1:50-200     IF(IHC-F) 1:50-200   IF(ICC) 1:50-200     Restrictions:   For Research Use only     Handling   Kester Comparison of the second	
UniProt:A6N7Y9Application DetailsApplication Notes:IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandling	
Application Details     Application Notes:   IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200     Restrictions:   For Research Use only     Handling   For Research Use only	
Application Notes:   IF(IHC-P) 1:50-200     IF(IHC-F) 1:50-200   IF(ICC) 1:50-200     Restrictions:   For Research Use only     Handling	
IF(IHC-F) 1:50-200   IF(ICC) 1:50-200   Restrictions: For Research Use only   Handling	
IF(ICC) 1:50-200   Restrictions:   For Research Use only   Handling	
Restrictions: For Research Use only Handling	
Handling	
Format: Liquid	
Concentration: 1 µg/µL	
Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proc	clin300 and
50 % Glycerol.	
Preservative: ProClin	
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which	should be
handled by trained staff only.	
Storage: -20 °C	
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date: 12 months	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN671638 | 03/06/2024 | Copyright antibodies-online. All rights reserved.