

## Datasheet for ABIN6719711

### WISP1 ELISA Kit

[Go to Product page](#)

#### Overview

Quantity:	1 kit
Target:	WISP1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	31.2 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

#### Product Details

Purpose:	Mouse WISP1 - CCN4 Sandwich ELISA Kit for Quantitative Detection
Brand:	AccuSignal™
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Production: Natural and recombinant mouse WISP1. There is no detectable cross-reactivity with other relevant proteins.
Sensitivity:	< 10 pg/mL
Components:	<ul style="list-style-type: none"><li>• Antibody-coated 96-well plate</li><li>• Target Protein Standard</li><li>• Detection antibody</li></ul>

## Product Details

- Detection reagent
- Diluent buffers
- Wash buffers
- Substrate Solution
- Stop solutions
- Adhesive covers

## Target Details

Target:	WISP1
Alternative Name:	WISP1/CCN4 ( <a href="#">WISP1 Products</a> )
Background:	<p>Synonyms: CCN family member 4, CCN4, WISP1, WISP1c, WISP1i, WISP1tc, Wnt 1 inducible signaling pathway protein 1, Wnt-1 induced secreted protein, WNT1 induced secreted protein 1, WNT1 inducible signaling pathway protein 1, WNT1 inducible-signaling pathway protein 1</p> <p>Background: WISP1(WNT1-Inducible Signaling Pathway Protein 1), also known as CCN4, is a matricellular protein that in humans is encoded by the WISP1 gene. WISP1 is induced by WNT1 and belongs to the CCN family, which includes connective tissue growth factor, cysteine-rich-61, and nephroblastoma overexpressed. By use of radiation hybrid mapping panels, the WISP1 was mapped to chromosome 8q24.1-q24.3, roughly 4 Mb distal to MYC. It found that 2 distinct systems demonstrated that WISP induction was associated with expression of WNT1. WISP1 genomic DNA was amplified in colon cancer cell lines and in human colon tumors, and its RNA was overexpressed in 84 % of tumors examined compared with patient-matched normal mucosa.</p>
Gene ID:	22402
NCBI Accession:	<a href="#">NP_061353</a>
UniProt:	<a href="#">O54775</a>
Pathways:	<a href="#">WNT Signaling</a> , <a href="#">Growth Factor Binding</a>

## Application Details

Application Notes:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 2000pg/mL,1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL mouse WISP1 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended
--------------------	--

Application Details

	that each mouse WISP1 standard solution and each sample be measured in duplicate.
Comment:	Standard: Expression system for standard: NSO, Immunogen sequence: T23-N367
Sample Volume:	100 µL
Plate:	Pre-coated
Restrictions:	For Research Use only

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiry Date:	12 months