

Datasheet for ABIN6719747

CXCL5 ELISA Kit**1** Image[Go to Product page](#)

Overview

Quantity:	1 kit
Target:	CXCL5
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.2 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin - citrate), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	<p>Synonyms: AMCF II, AMCFII, C X C motif chemokine 5, Chemokine (C X C motif) ligand 5, CXCL 5, ENA 78, ENA78, Epithelial derived neutrophil activating protein 78, GCP 2, LIX Neutrophil activating peptide ENA 78, Neutrophil activating protein 78, SCYB5, Scyb6, Small inducible cytokine B5, Small inducible cytokine subfamily B (Cys X Cys) member 5 (epithelial derived neutrophil activating peptide 78), Small inducible cytokine subfamily B, member 15</p> <p>Background: C-X-C motif chemokine 5 is a protein that in humans encoded by the CXCL5 gene. The protein encoded by this gene, CXCL5 is a small cytokine belonging to the CXC chemokine family that is also known as epithelial-derived neutrophil-activating peptide 78(ENA-78). It is</p>

Product Details

produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon IFN-gamma. This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2. The gene for CXCL5 is encoded on four exons and is located on human chromosome 4 amongst several other CXC chemokine genes. CXCL5 has been implicated in connective tissue remodeling. CXCL5 plays a role in reducing sensitivity to sunburn pain in some subjects, and is a potential target which can be utilized to understand more about pain in other inflammatory conditions like arthritis and cystitis.

Gene Name: CXCL5

Production: Natural and recombinant human CXCL5. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: E.coli, Immunogen sequence: A37-N114

Target Details

Target:	CXCL5
Alternative Name:	CXCL5 - ENA-78 (CXCL5 Products)
Gene ID:	6374
NCBI Accession:	NP_002985
UniProt:	P42830
Pathways:	Cellular Response to Molecule of Bacterial Origin , Regulation of Leukocyte Mediated Immunity

Application Details

Application Notes:	<p>Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL human CXCL5 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma (heparin, EDTA, citrate) to each empty well. It is recommended that each human CXCL5 standard solution and each sample be measured in duplicate.</p> <p>Blood Product Anticoagulant: Heparin Sodium</p> <p>ELISA Dilution: 31.2pg/mL-2000pg/mL</p>
Sample Volume:	100 µL


Application Details

Plate:	Pre-coated
Restrictions:	For Research Use only

Handling

Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images



ELISA

Image 1. Human CXCL5/ENA-78 Accusignal ELISA Kit

Human CXCL5/ENA-78 AccuSignal ELISA Kit standard curve.

Assay Range: 31.2pg/ml-2000pg/ml.

Sensitivity: <10pg/ml.