

Datasheet for ABIN6719756 **VEGF ELISA Kit**



[Go to Product page](#)

1 Image

Overview

Quantity:	1 kit
Target:	VEGF
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.2 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin - citrate), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	<p>Synonyms: MGC70609, MVCD1, Vascular endothelial growth factor, Vascular endothelial growth factor A, Vascular permeability factor, VEGF A, VEGFA, VPF</p> <p>Background: Vascular permeability factor/vascular endothelial growth factor(VPF/VEGF), a potent cytokine expressed by most malignant tumors, has critical roles in vasculogenesis and both physiological and pathological angiogenesis. VEGF produced by tumor cells potentially stimulates endothelial cell proliferation and angiogenesis and plays a key role in the pathophysiology of several neoplasias. VEGF may also play a pivotal role in mediating the development and progression of diabetic retinopathy. VEGF, a major regulator of angiogenesis,</p>

Product Details

binds to two receptor tyrosine kinases, KDR/Flk-1 and Flt-1. The VEGF gene is mapped by fluorescence in situ hybridization to chromosome 6p12.

Gene Name: VEGFA

Production: Natural and recombinant human VEGF. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: sf21, Immunogen sequence: A27-R191

Target Details

Target: VEGF

Alternative Name: VEGF ([VEGF Products](#))

Gene ID: 7422

NCBI Accession: [NP_001020537](#)

UniProt: [P15692](#)

Application Details

Application Notes: Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL human VEGF standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma (heparin, EDTA, citrate) to each empty well. We recommend that each human VEGF standard solution and each sample is measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 31.2pg/mL-2000pg/mL(humanserum,plasma)15.6pg/mL-

1000pg/mL(cellculturesupernates)

Other Performance Data: DOE: 2-8° C - 05/14/18 -20° C - 11/14/18

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only


Handling

Storage: RT,4 °C,-20 °C

Handling

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images



ELISA

Image 1. Human VEGF Accusignal ELISA Kit Human VEGF Accusignal ELISA Kit standard curve. Assay Range: 31.2pg/ml-2000pg/ml(humanserum,plasma)15.6pg/ml-1000pg/ml(cellculturesupernates). Sensitivity: <1pg/ml.