

Datasheet for ABIN6719781

E-cadherin ELISA Kit

1 Image



Go to Product page

Overview

Quantity:	1 kit
Target:	E-cadherin (CDH1)
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	156 pg/mL - 10000 pg/mL
Minimum Detection Limit:	156 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen Cell Culture Supernatant, Plasma (heparin), Serum	
Sample Type:		
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Characteristics:	Synonyms: Arc 1, Cadherin 1, cadherin 1 type 1 E-cadherin, Cadherin1, CAM 120/80, CD 234,	
	CD324, CD324 antigen, CDH1, CDHE, ECAD, Epithelial cadherin, epithelial calcium dependant	
	adhesion protein, LCAM, Liver cell adhesion molecule, UVO, Uvomorulin	
	Background: E-Cadherin, also called Cadherin 1(CDH1), Uvomorulin or calcium-dependent	
	ashesion protein, epithelial. E-cadherin is a Ca(2+)-dependent epithelial cell-cell adhesion	
	molecule. Downregulation of E-cadherin expression often correlates with strong invasive	
	potential and poor prognosis of human carcinomas. The gene spans a region of approximately	
	100 kb, and its location on chromosome 16q22.1. It contains 16 exons and a 65-kb-long intron	

2. E-cadherin gene mutations may contribute to the development of diffusely growing gastric carcinomas. E-cadherin plays a central part in the process of epithelial morphogenesis and acts as a strong invasion suppressor in experimental tumor cell systems.

Gene Name: CDH1

Production: Natural and recombinant mouse E-Cadherin. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: D157-V709

Target Details

Target:	E-cadherin (CDH1)	
Alternative Name:	E-Cadherin (CDH1 Products)	
Gene ID:	12550	
NCBI Accession:	NP_033994	
UniProt:	P09803	
Pathways:	WNT Signaling, Sensory Perception of Sound, Cell-Cell Junction Organization, Tube Formation	

Application Details

Δnn	lication	Notae.

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 10000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 313pg/mL, 156pg/mL mouse E-Cadherin standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin) to each empty well. We recommend that each mouse E-Cadherin standard solution and each sample is measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 156pg/mL-10000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only


Handling

Storage: RT,4 °C,-20 °C

Storage Comment:

Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images



ELISA

Image 1. Mouse E-Cadherin Accusignal ELISA Kit Mouse E-Cadherin AccuSignal ELISA Kit standard curve. Assay Range: 156pg/ml-10000pg/ml. Sensitivity: <15pg/ml.