

Datasheet for ABIN6719784

Relaxin 1 ELISA Kit



Overview

Quantity:	1 kit
Target:	Relaxin 1 (RLN1)
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	78 pg/mL - 5000 pg/mL
Minimum Detection Limit:	78 pg/mL
Application:	ELISA

Product Details

Purpose:	Mouse Relaxin 1 Sandwich ELISA Kit for Quantitative Detection
Brand:	AccuSignal™
Sample Type:	Cell Culture Supernatant, Plasma (heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Production: Natural and recombinant mouse Relaxin 1. There is no detectable cross-reactivity with other relevant proteins.
Sensitivity:	< 10 pg/mL
Components:	Antibody-coated 96-well plateTarget Protein Standard

· Detection antibody

- Detection reagent
- · Diluent buffers
- Wash buffers
- · Substrate Solution
- · Stop solutions
- · Adhesive covers

Target Details

Target:	Relaxin 1 (RLN1)
Alternative Name:	Relaxin 1 (RLN1 Products)
Background:	Synonyms: H1RLX, Preprorelaxin H1, PreprorelaxinH1, Prorelaxin, Prorelaxin H1, ProrelaxinH1, REL1_HUMAN, Relaxin 1 (H1), Relaxin A chain, Relaxin H1, Relaxin1, RelaxinH1, RLN 1, RLN1, RLX H1, RLXH 1, RLXH Background: Relaxin 1 is a member of relaxin-like peptide family. Relaxin gene maps to mouse chromosome 19 near D19Mit23. Relaxin is a peptide hormone produced by the corpora lutea of ovaries during pregnancy in many mammalian species, including man. Relaxin widens the pubic bone and facilitates labor, it also softens the cervix(cervical ripening), and relaxes the uterine musculature. However, its significance may reach much further. Relaxin affects collagen metabolism, inhibiting collagen synthesis and enhancing its breakdown by increasing matrix metalloproteinases. It also enhances angiogenesis and is a potent renal vasodilator.
Gene ID:	19773
NCBI Accession:	NP_035402
UniProt:	P47932

Application Details

Application Notes:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of
	5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 313pg/mL, 156pg/mL, 78pg/mL mouse
	Relaxin 1 standard solutions into the pre-coated 96-well plate. Add 0.1 mL of the sample diluent
	buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse
	cell culture supernates, serum or plasma(heparin) to each empty well. We recommend that
	each mouse Relaxin 1 standard solution and each sample is measured in duplicate.
Comment:	Standard: Expression system for standard: E.coli, Immunogen sequence: R23-R54(B)&E161-C185(A)

Application Details

Sample Volume:	100 μL
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiry Date:	12 months