

Datasheet for ABIN6719819 **CST3 ELISA Kit**



[Go to Product page](#)

1 Image

Overview

Quantity:	1 kit
Target:	CST3
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	312 pg/mL - 20000 pg/mL
Minimum Detection Limit:	312 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	<p>Synonyms: ARMD11, bA218C14.4 (cystatin C), Cst3, Cystatin-3, Cystatin-C, CYTC_HUMAN, Gamma-trace, Neuroendocrine basic polypeptide, Post-gamma-globulin</p> <p>Background: Cystatin C or cystatin 3(formerly gamma trace, post-gamma-globulin or neuroendocrine basic polypeptide), a protein encoded by the CST3 gene, was originally described as a constituent of normal cerebrospinal fluid(CSF) and of urine from patients with renal failure. Cystatin 3 has a low molecular weight(approximately 13.3 kilodaltons), and it is removed from the bloodstream by glomerular filtration in the kidneys. In mouses, all cells with a nucleus(cell core containing the DNA) produce cystatin C as a chain of 120 amino acids. It is</p>

Product Details

found in virtually all tissues and bodily fluids. Cystatin C, which belongs to the type II cystatin gene family, is a potent inhibitor of lysosomal proteinases (enzymes from a special subunit of the cell that break down proteins) and probably one of the most important extracellular inhibitors of cysteine proteases (it prevents the breakdown of proteins outside the cell by a specific type of protein degrading enzymes). Moreover, cystatin C is involved in network reorganization in the ...

Gene Name: CST3

Production: Natural and recombinant mouse Cystatin C. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: M1-A140

Target Details

Target: CST3

Alternative Name: Cystatin C ([CST3 Products](#))

Gene ID: 1471

NCBI Accession: [NP_000090](#)

UniProt: [P01034](#)

Application Details

Application Notes: Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 20000pg/mL, 10000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 312pg/mL mouse Cystatin C standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum, plasma (heparin, EDTA) or urine to each empty well. It is recommended that each mouse Cystatin C standard solution and each sample be measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 312pg/mL-20000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated


Restrictions: For Research Use only

Handling

Storage: RT,4 °C,-20 °C

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images



ELISA

Image 1. Mouse Cystatin C Accusignal ELISA Kit Mouse Cystatin C AccuSignal ELISA Kit standard curve. Assay Range: 312pg/ml-20000pg/ml. Sensitivity: <10pg/ml.