

Datasheet for ABIN6719893

ERBB3 ELISA Kit



Overview

Quantity:	1 kit
Target:	ERBB3
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	156 pg/mL - 10000 pg/mL
Minimum Detection Limit:	156 pg/mL
Application:	ELISA

Product Details

Purpose:	Human ERBB3 - Her3 Sandwich ELISA Kit for Quantitative Detection
Brand:	AccuSignal™
Sample Type:	Cell Culture Lysate, Cell Culture Supernatant, Plasma (EDTA - heparin), Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Production: Natural and recombinant human ERBB3. There is no detectable cross-reactivity with other relevant proteins.
Sensitivity:	< 10 pg/mL
Components:	Antibody-coated 96-well plateTarget Protein Standard

- Detection antibody
- · Detection reagent
- · Diluent buffers
- · Wash buffers
- · Substrate Solution
- · Stop solutions
- · Adhesive covers

Target Details

Target:	ERBB3
Alternative Name:	ERBB3/Her3 (ERBB3 Products)
Background:	Synonyms: c erbB3, Erb b2 receptor tyrosine kinase 3, ERBB3 protein, erbB3 S, ERBB3_HUMAN
	Glial growth factor receptor, HER 3, R3, Human epidermal growth factor receptor 3, LCCS2,
	MDA BF 1, p180 ErbB3, p45 sErbB3, p85 sErbB3, proto-oncogene-like protein c ErbB3, Recepto
	tyrosine protein kinase erbB 3, Tyrosine kinase type cell surface receptor HER3, Tyrosine
	kinase-type cell surface receptor HER3, v erb b2 avian erythroblastic leukemia viral oncogene
	homolog 3, v erb b2 erythroblastic leukemia viral oncogene homolog 3
	Background: Receptor tyrosine-protein kinase erbB-3, also known as HER3 (human epidermal
	growth factor receptor 3), is a membrane bound protein that in humans is encoded by the
	ERBB3 gene. ErbB3 has been shown to bind the ligands heregulin and NRG-2. Ligand binding
	causes a change in conformation that allows for dimerization, phosphorylation, and activation
	of signal transduction. ErbB3 can heterodimerize with any of the other three ErbB family
	members. The theoretical ErbB3 homodimer would be non-functional because the kinase-
	impaired protein requires transphosporylation by its binding partner to be active. Unlike the
	other ErbB receptor tyrosine kinase family members which are activated through
	autophosphorylation upon ligand binding, ErbB3 is found to be kinase impaired, having only
	1/1000th the autophosphorylation activity of EGFR and no ability to phosphorylate other
	proteins. Therefore, ErbB3 must act as anallosteric activator.
Gene ID:	2065
NCBI Accession:	NP_001005915
UniProt:	P21860
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway

Application Details

Application Notes:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the
	10,000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 312pg/mL, 156pg/mL human
	ERBB3 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent
	buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human
	cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended
	that each human ERBB3 standard solution and each sample be measured in duplicate.
Comment:	Standard: Expression system for standard: NSO, Immunogen sequence: S20-T643
Sample Volume:	100 μL
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid
	cycles of freezing and thawing.
Expiry Date:	12 months