


Datasheet for ABIN6719912

MMP2 ELISA Kit





Overview

Quantity:	1 kit
Target:	MMP2
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	625 pg/mL - 40000 pg/mL
Minimum Detection Limit:	625 pg/mL
Application:	ELISA

Product Details

Purpose:	Mouse MMP-2 Sandwich ELISA Kit for Quantitative Detection
Brand:	AccuSignal™
Sample Type:	Cell Culture Supernatant, Plasma (heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Production: Natural and recombinant mouse total MMP-2. There is no detectable cross-reactivity with other relevant proteins.
Sensitivity:	< 10 pg/mL
Components:	Antibody-coated 96-well plateTarget Protein Standard

Detection antibody

- Detection reagent
- · Diluent buffers
- Wash buffers
- · Substrate Solution
- Stop solutions
- · Adhesive covers

Target Details

Target:	MMP2
Alternative Name:	MMP-2 (MMP2 Products)
Background:	Synonyms: 73 kDa gelatinase, 72 kDa type IV collagenase, CLG 4, CLG 4A, CLG4, CLG4A,
	Collagenase type IV A, EC 3.4.24.24, Gelatinase A, Gelatinase alpha, Gelatinase neutrophil,
	Matrix metallopeptidase 2 (gelatinase A, 72 kDa gelatinase, 72 kDa type IV collagenase)
	Background: Type IV collagenase, 72-kD, is officially designated matrix metalloproteinase-
	2(MMP2). It is also known as gelatinase, 72-kD. MMP-2 plays an essential role in angiogenesis
	and arteriogenesis, two processes critical to restoration of tissue perfusion after ischemia.
	MMP-2 expression is increased in tissue ischemia, but the responsible mechanisms remain
	unknown. Matrix metalloproteinases(MMPs) catalyze extracellular matrix degradation. Control
	of their activity is a promising target for therapy of diseases characterized by abnormal
	connective tissue turnover. MMPs are expressed as latent proenzymes that are activated by
	proteolytic cleavage that triggers a conformational change in the propeptide(cysteine switch).
	The structure of proMMP-2 reveals how the propeptide shields the catalytic cleft and that the
	cysteine switch may operate through cleavage of loops essential for propeptide stability. The
	gene is localized to 16q21 using somatic cell hybrids and in situ hybridization. The standard
	product used in this kit is recombinant mouse MMP-2, consisting of 662 amino acids with the
	molecular mass of 72KDa. The detected MMP-2 includes zymogen and active enzyme.
Gene ID:	17390
NCBI Accession:	NP_032636
UniProt:	P33434
Pathways:	Activation of Innate immune Response
Application Details	
Application Notes:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the

40000pg/mL, 20,000pg/mL,10,000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL mouse MMP-2 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma (heparin) to each empty well. It is recommended that each mouse MMP-2 standard solution and each sample be measured in duplicate.

Comment:

Standard: Expression system for standard: NSO, Immunogen sequence: A30-C662

Sample Volume:

100 µL

Plate:

Pre-coated

Restrictions:

For Research Use only

Handling

Storage:

4 °C,-20 °C


Storage Comment:

Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Expiry Date:

12 months

Images



ELISA

Image 1. Mouse MMP-2 Accusignal ELISA Kit Mouse MMP-2 AccuSignal ELISA Kit standard curve. Assay Range: 625pg/ml-40000pg/ml. Sensitivity: <10pg/ml.