

# Datasheet for ABIN6719943

# **RANKL ELISA Kit**





#### Go to Product page

#### Overview

Quantity:	1 kit
Target:	RANKL (TNFSF11)
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	93.7 pg/mL - 6000 pg/mL
Minimum Detection Limit:	93.7 pg/mL
Application:	ELISA

## **Product Details**

Product Details	
Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	Synonyms: MGC29565, MGC93568, OCIF, OPG, Osteoclastogenesis inhibitory factor,
	Osteoprotegerin, TNFRSF 11B, TNFRSF11B, TR 1, TR1, TR11B_HUMAN, Tumor necrosis factor
	receptor superfamily member 11B
	Background: Osteoprotegerin(OPG) is identical to osteoclastogenesis inhibitory factor(OCIF), a
	soluble member of the tumor-necrosis factor receptor family that inhibits osteoclastogenesis.
	OPG is considered to play an important role in the regulation of bone resorption by modifying
	osteoclast differentiation. Osteoprotegerin is an independent risk factor for the progression of
	atherosclerosis and onset of cardiovascular disease. It can act as a soluble factor in the

regulation of bone mass and imply a utility for OPG in the treatment of osteoporosis associated with increased osteoclast activity. OPG system may play a critical role in the development of osteolytic bone disease in multiple myeloma and that targeting this system may have therapeutic potential. OPG also plays a role in the vascular system.

Gene Name: TNFRSF11B

Production: Natural and recombinant mouse OPG. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: E22-L401

#### **Target Details**


Target:	RANKL (TNFSF11)
Alternative Name:	OPG (TNFSF11 Products)
Gene ID:	18383
NCBI Accession:	NP_032790
UniProt:	008712
Pathways:	NF-kappaB Signaling

Application Details	
Application Notes:	Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot
	0.1 mL per well of the 6000pg/mL, 3000pg/mL, 1500pg/mL, 750pg/mL, 419pg/mL,
	187.5pg/mL, 93.7pg/m mouse OPG standard solutions into the precoated 96-well plate. Add
	0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each
	properly diluted sample of mouse cell culture supernates, serum or plasma(EDTA) to each
	empty well. It is recommended that each mouse OPG standard solution and each sample be
	measured in duplicate.
	Blood Product Anticoagulant: Sodium EDTA
	ELISA Dilution: 93.7pg/mL-6000pg/mL
Sample Volume:	100 μL
Plate:	Pre-coated
Restrictions:	For Research Use only

## Handling

Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid
	cycles of freezing and thawing.

## **Images**



#### **ELISA**

**Image 1.** Mouse OPG Accusignal ELISA Kit Mouse OPG AccuSignal ELISA Kit standard curve. Assay Range: 93.7pg/ml-6000pg/ml. Sensitivity: <5pg/ml.