antibodies -online.com





TWEAK ELISA Kit





Overview

Quantity:	1 kit
Target:	TWEAK (TNFSF12)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	62.5 pg/mL - 4000 pg/mL
Minimum Detection Limit:	62.5 pg/mL
Application:	ELISA

Product Details	
Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	Synonyms: APO 3 ligand, APO 3L, APO3 ligand, APO3/DR3 ligand, APO3L, CD255, DR3LG,
	MGC129581, MGC20669, TNF related weak inducer of apoptosis, TNF-related weak inducer of
	apoptosis, TNF12_HUMAN, TNFRS12a, TNFSF 12, TNFSF12, TNFSF12 protein, Tumor necrosis
	factor (ligand) superfamily member 12, Tumor necrosis factor ligand superfamily member 12,
	Tumor necrosis factor superfamily member 12, TWEAK, UNQ181/PRO207
	Background: Tumor necrosis factor ligand superfamily member 12 also known as TNF-related
	weak inducer of apoptosis(TWEAK) is a protein that in humans is encoded by the TNFSF12
	gene. The protein encoded by this gene is a cytokine that belongs to the tumor necrosis

factor(TNF) ligand family. This protein is a ligand for the FN14/TWEAKR receptor. This cytokine has overlapping signaling functions with TNF, but displays a much wider tissue distribution. This cytokine can induce apoptosis via multiple pathways of cell death in a cell type-specific manner. This cytokine is also found to promote proliferation and migration of endothelial cells, and thus acts as a regulator ofangiogenesis. The TNFSF12 gene lies 878 bp upstream of the putative transcriptional start site of the TNFSF13 gene on chromosome 17p13.1.

Gene Name: TNFSF12

Production: Natural and recombinant human TNFSF12. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: E.coli, Immunogen sequence: S94-H249

Target Details


Target:	TWEAK (TNFSF12)
Alternative Name:	TNFSF12 (TNFSF12 Products)
Gene ID:	407977
NCBI Accession:	NP_003800
UniProt:	043508
Pathways:	Apoptosis

Application Details	
Application Notes:	Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot
	0.1 mL per well of the 4000pg/mL, 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL,
	62.5pg/mL human TNFSF12 standard solutions into the precoated 96-well plate. Add 0.1 mL of
	the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted
	sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well.
	It is recommended that each human TNFSF12 standard solution and each sample be
	measured in duplicate.
	Blood Product Anticoagulant: Heparin Sodium
	ELISA Dilution: 62.5pg/mL-4000pg/mL
Sample Volume:	100 μL
Plate:	Pre-coated
Restrictions:	For Research Use only

Handling

Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid
	cycles of freezing and thawing.

Images



ELISA

Image 1. Human TNFSF12 Accusignal ELISA Kit Human TNFSF12 AccuSignal ELISA Kit standard curve. Assay Range: 62.5pg/ml-4000pg/ml. Sensitivity: <10pg/ml.