antibodies .- online.com





FASL ELISA Kit





Overview

| Quantity: | 1 kit |
|--------------------------|-------------------------|
| Target: | FASL |
| Reactivity: | Human |
| Method Type: | Sandwich ELISA |
| Detection Range: | 15.6 pg/mL - 1000 pg/mL |
| Minimum Detection Limit: | 15.6 pg/mL |
| Application: | ELISA |

Product Details

| Purpose: | Sandwich ELISA for Quantitative Detection of Antigen |
|--------------------|---|
| Sample Type: | Cell Culture Supernatant, Plasma (EDTA - heparin - citrate), Serum |
| Analytical Method: | Quantitative |
| Detection Method: | Colorimetric |
| Characteristics: | Synonyms: Apoptosis (APO 1) antigen ligand 1, Apoptosis antigen ligand, APT1LG1, APTL, |
| | CD178, CD95 L, CD95 ligand, CD95L, CD95L protein, Fas antigen ligand, FASL, FASLG, TNF |
| | superfamily member 6, TNFSF6, Tumor necrosis factor (ligand) superfamily member 6, Tumor |
| | necrosis factor ligand superfamily member 6 |
| | Background: Fas ligand(FasL or CD95L) is a type-II transmembrane protein that belongs to the |
| | tumor necrosis factor(TNF) family. Its binding with its receptor induces apoptosis. The human |

FASL gene consists of approximately 8 kb and is split into 4 exons. Fas ligand/receptor

interactions play an important role in the regulation of the immune system and the progression

of cancer. Fas ligand or FasL is a homotrimeric type II transmembrane protein. It signals through trimerization of FasR, which spans the membrane of the "target" cell. This trimerization usually leads to apoptosis, or cell death. Soluble Fas ligand is generated by cleaving membrane-bound FasL at a conserved cleavage site by the external matrix metalloproteinase MMP-7.

Gene Name: FASLG

Production: Natural and recombinant human FASL. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: CHO, Immunogen sequence: P134-L281

Target Details


| Target: | FASL |
|-------------------|---|
| Alternative Name: | FASL (FASL Products) |
| Gene ID: | 14103 |
| NCBI Accession: | NP_001192172 |
| UniProt: | Q99PH8 |
| Pathways: | Apoptosis, EGFR Signaling Pathway, Production of Molecular Mediator of Immune Response, |
| | Positive Regulation of Endopeptidase Activity |

| Application Details | |
|---------------------|--|
| Application Notes: | Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot |
| | 0.1 mL per well of the 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL, |
| | 15.6pg/mL human FASL standard solutions into the precoated 96-well plate. Add 0.1 mL of the |
| | sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted |
| | sample of human cell culture supernates, serum or plasma(heparin, EDTA, citrate) to each |
| | empty well. It is recommended that each human FASL standard solution and each sample be |
| | measured in duplicate. |
| | Blood Product Anticoagulant: Heparin Sodium |
| | ELISA Dilution: 15.6pg/mL-1000pg/mL |
| Sample Volume: | 100 μL |
| Plate: | Pre-coated |
| Restrictions: | For Research Use only |

Handling

| Storage: | RT,4 °C,-20 °C |
|------------------|--|
| Storage Comment: | Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only |
| | prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing. |
| | oyoled of freezing and thawing. |

Images



ELISA

Image 1. Human FASL Accusignal ELISA Kit Human FASL AccuSignal ELISA Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <2pg/ml.