
antibodies -online.com





VEGF ELISA Kit





Go to Product page

_						
0	V	0	r٧	/[Θ	M

Quantity:	1 kit
Target:	VEGF
Reactivity:	Rat
Method Type:	Sandwich ELISA
Detection Range:	15.6 pg/mL - 1000 pg/mL
Minimum Detection Limit:	15.6 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen		
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin - citrate), Serum		
Analytical Method:	Quantitative		
Detection Method:	Colorimetric		
Characteristics:	Synonyms: MGC70609, MVCD1, Vascular endothelial growth factor, Vascular endothelial growth factor A, Vascular permeability factor, VEGF A, VEGFA, VPF		

Background: Vascular permeability factor, VEGF A, VEGFA, VPF
Background: Vascular permeability factor/vascular endothelial growth factor(VPF/VEGF), a
potent cytokine expressed by most malignant tumors, has critical roles in vasculogenesis and
both physiological and pathological angiogenesis. VEGF produced by tumor cells potently
stimulates endothelial cell proliferation and angiogenesis and plays a key role in the
pathophysiology of several neoplasias. VEGF may also play a pivotal role in mediating the
development and progression of diabetic retinopathy. VEGF, a major regulator of angiogenesis,

binds to two receptor tyrosine kinases, KDR/Flk-1 and Flt-1. The VEGF gene is mapped by fluorescence in situ hybridization to chromosome 6p12. The standard product used in this kit is recombinant rat VEGF164, which is a 25KDa single chain as well as a 50KDa dimer.

Gene Name: VEGFA

Production: Natural and recombinant rat VEGF. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: A27-R190

Endotoxin level: Low endotoxin

Target Details

Target:	VEGF
Alternative Name:	VEGF (VEGF Products)
Gene ID:	83785
NCBI Accession:	NP_001103803
UniProt:	B5DEK7

Application Details

Application Notes:

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.3pg/mL, 15.6pg/mL rat VEGF standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of rat cell culture supernates, serum or plasma(heparin, EDTA, citrate) to each empty well. We recommend that each rat VEGF standard solution and each sample is measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 15.6pg/mL-1000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only

Handling


Storage: RT,4 °C,-20 °C

Handling

Storage Comment:

Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images



ELISA

Image 1. Rat VEGF Accusignal ELISA Kit Rat VEGF AccuSignal ELISA Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <1pg/ml.